Neural precursor and stem cells

Patent number:

EP1529838

Publication date:

2005-05-11

Inventor:

BOSIO ANDREAS DR (DE); CREMER HAROLD (FR);

PENNARTZ SANDRA (DE)

Applicant:

MEMOREC BIOTEC GMBH (DE)

Classification:

- international:

C12N5/06; G01N33/53

- european:

C12N5/06B8P; G01N33/68V2

Application number: EP20030025506 20031106

Priority number(s): EP20030025506 20031106

Cited documents:

EP1354943 XP00227572 XP00222350

XP00103392

XP00902890

Report a data error he

Abstract of EP1529838

A cell population comprising at least 5% neural stem cells, the stem cells being characterized by an expression of ASCT2 or KIAA0152, is new. - Independent claims are also included for the following: - (1) a method for isolating the cell population cited above; - (2) a medicament comprising the above cell population; and - (3) a monoclonal antiboc directed against ASCT2. - ACTIVITY - Neuroprotective; Nootropic; Antiparkinsonian; Cerebroprotective; Vasotropic; No biological data given. - MECHANISM OF ACTION - Cell Therapy.

Data supplied from the esp@cenet database - Worldwide



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11) EP 1 529 838 A1

(12)

EUROPÄISCHE PATENTANMELDUNG

- (43) Veröffentlichungstag: 11.05.2005 Patentblatt 2005/19
- (51) Int Cl.7: C12N 5/06, G01N 33/53

- (21) Anmeldenummer: 03025506.1
- (22) Anmeldetag: 06.11.2003
- (84) Benannte Vertragsstaaten:
 AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
 HU IE IT LI LU MC NL PT RO SE SI SK TR
 Benannte Erstreckungsstaaten:
 AL LT LV MK
- (71) Anmelder: Memorec Biotec GmbH 50829 Köln (DE)
- (72) Erfinder:
 - Bosio, Andreas, Dr. 50931 Köln (DE)

- Cremer, Harold 13400 Marseille (FR)
 Pennartz, Sandra 50670 Köln (DE)
- (74) Vertreter: Schreiber, Christoph, Dr. et al Patentanwälte von Kreisler Selting Werner, Postfach 10 22 41 50462 Köln (DE)
- (54) Neurale Vorläufer- und Stammzellen
- (57) Zellpopulation, dadurch gekennzeichnet, dass mindestens 5% der Zellen neurale Vorläuferzellen sind, die wenigstens einen der in **Liste A** oder **Liste B** aufgeführten Marker aufweisen.

2

Beschreibung

[0001] Die vorliegende Erfindung betrifft Zellpopulationen von neuralen Vorläuferzellen bzw. neuralen Stammzellen sowie Verfahren zur Isolierung entsprechender Zellen.

1

[0002] Der Ausgangspunkt für die Generierung der über tausend verschiedenen neuronalen und glialen Zelltypen des Nervensystems von Vertebraten sind multipotente, neurale Stammzellen des embryonalen Neuroepitheliums (Williams, B. P., Read, J. & Price, J. (1991): The generation of neurons and oligodendrocytes from a common precursor cell. Neuron 7(4), 685-93), (Davis, A. A. & Temple, S. (1994): A self-renewing multipotential stem cell in embryonic rat cerebral cortex. Nature 372(6503), 263-6), (Weiss, S., Dunne, C., Hewson, J., Wohl, C., Wheatley, M., Peterson, A. C. & Reynolds, B. A. (1996): Multipotent CNS stem cells are present in the adult mammalian spinal cord and ventricular neuroaxis. J Neurosci 16(23), 7599-609).

[0003] In den vergangenen Jahren wurde durch verschiedene Arbeitsgruppen gezeigt, dass solche sich selbst erneuernden, multipotenten Vorläuferzellen nicht nur während der Entwicklung, sondern auch im adulten Gehirn zu finden sind (Gage, F. H. (2000): Mammalian neural stem cells. *Science* 287(5457), 1433-8). Vor allem um die lateralen Ventrikel des Vorderhirns findet die Bildung von neuralen Vorläuferzellen lebenslang statt. Diese wandern hauptsächlich, wenn auch nicht exklusiv, in den Bulbus olfaktorius, um dort in GABA-erge Interneurone zu differenzieren.

[0004] Über die genaue Lokalisation der multipotenten Stammzellen, die dieser sekundären Neurogenese zugrunde liegen, wird derzeit noch spekuliert: Johansson et al. beschrieben ependymale Zellen entlang des Lumen der adulten, ventrikulären Zone mit den Eigenschaften multipotenter Stammzellen (Johansson, C. B., Svensson, M., Wallstedt, L., Janson, A. M. & Frisen, J. (1999b): Neural stem cells in the adult human brain. Exp. Cell Res 253(2), 733-6), während Doetsch et al. Astrocyten der subventrikulären Zone als multipotente Stammzellen identifizierten (Doetsch, F., Caille, I., Lim, D. A., Garcia-Verdugo, J. M. & Alvarez-Buylla, A. (1999): Subventricular zone astrocytes are neural stem cells in the adult mammalian brain. Cell 97(6), 703-16). Eine absolut eindeutige Identifizierung dieser adulten Stammzellen in vivo ist jedoch bis heute, hauptsächlich mangels geeigneter Marker, nicht gelungen.

[0005] Neben ihrer Bedeutung im olfaktorischen System ist das therapeutische Potential der adulten Stammzellen von besonderem Interesse. Aufgrund ihrer Multipotenz weisen neurale Stammzellen bemerkenswerte Formbarkeit auf und könnten daher durch Zusatz von verschiedenen Faktoren zur Erzeugung verschiedener Neuronentypen eingesetzt werden. Die anschließende Transplantation der so entwickelten spezialisierten Zellen könnte zur Behandlung von neurologischen Krankheiten Alzheimer, Parkinson, Folgen von

Schädel-Hirn-Traumata und Schlaganfall beitragen. Voraussetzung dafür ist die Charakterisierung der verschiedenen, neuralen Differenzierungsstufen sowie die Identifizierung der Faktoren, die die Differenzierungsprogramme der Stammzellen steuern. Gegenüber den embryonalen Stammzellen haben die adulten den Vorteil, dass sie erstens keine abstoßende Immunreaktion auslösen würden, weil sie dem Körper des Patienten entstammen, folglich ihre Transplantation ohne Immunsuppression erfolgen könnte, und zweitens ihre Gewinnung ethisch unbedenklich ist.

[0006] Die Erforschung der Eigenschaften neuraler Stammzellen und embronaler Stammzellen des Menschen ist aus ethischen Aspekten praktisch nicht oder nur sehr eingeschränkt möglich. Daher wurden alle explorativen Arbeiten ausgehend von Mäusen und Mauszellen durchgeführt. Wie bereits beschrieben war die Isolierung von neuralen Stammzellen bisher nicht möglich, da dieser Zelltyp nicht eindeutig charakterisiert war und keine geeigneten Marker zur Identifizierung und Anreicherung zur Verfügung standen.

[0007] Aufgabe der vorliegenden Erfindung war es daher Verfahren zu entwickeln, die eine Isolation von neuralen Vorläuferzellen und neuralen Stammzellen erlauben und entsprechende Zellpopulation, enthaltend diese Zelle bereitzustellen.

[0008] Erfindungsgemäß wird die Aufgabe gelöst durch die Identifizierung von Markern, die entsprechende Zellen aufweisen.

[0009] Marker ist ein Gen, das mit Hilfe der Serial Analysis of Genexpression (SAGE) in entsprechenden Zellen gefunden wird.

[0010] Methodisch beruht SAGE auf der Isolierung von 14 bp großen DNA Fragmenten (Tags), die jeweils charakteristisch für eine mRNA-Spezies sind. Die Tags, repräsentativ für alle in der zu untersuchenden Zelle vorliegenden mRNA Moleküle, werden zu langen Polymeren verbunden, die im letzten Schritt der Methode sequenziert werden. Die Frequenz, mit der ein Tag sequenziert wird, ist direkt proportional zur Kopienzahl der mRNA-Moleküle im untersuchten Ausgangsmaterial (Velculescu, V. E., Zhang, L., Vogelstein, B. & Kinzler, K. W. (1995): Serial analysis of gene expression. Science 270(5235), 484-7). Durch die computerunterstützte Auswertung der Sequenzdaten entsteht ein digitales Expressionsprofil, das beliebig oft und ohne zusätzliche Laborarbeit mit Expressionsprofilen anderer Gewebe verglichen werden kann (Meta-Analyse).

[0011] Den so identifizierten Gene sind eindeutigen Nummern zugeordnet, die beispielsweise als SAGEmap von National Center for Biotechnology Information (NCBI) bereitgestellt werden (www.ncbi.nlm.nin.gov/SAGE).

[0012] Gegenstand der Erfindung sind zum einen Zellpopulationen, bei denen mindestens 5% der Zellen neurale Vorläuferzellen sind, die wenigstens einen der in Liste A oder Liste B aufgeführten Marker aufweisen.
[0013] Bevorzugt weisen entsprechende neurale Vor-

10

läuferzellen wenigstens zwei, drei, vier oder fünf der in Liste A oder B aufgeführten Marker auf.

3

[0014] In bevorzugten Ausführungsformen weisen entsprechende neurale Vorläuferzellen keinen der in Liste C aufgeführten Marker auf.

[0015] Bevorzugt ist der Gehalt an neuralen Vorläuferzellen in der Zellpopulation hoch, d.h. mindestens 10%, bevorzugt mindestens 25%, noch mehr bevorzugt mehr als 50% und am meisten bevorzugt über 90%.

[0016] Entsprechende neurale Vorläuferzellen sind vorzugsweise aus Hirngewebe erhältlich.

[0017] In einer Ausführungsform handelt es sich dabei um eine murine Zellpopulation.

[0018] Gegenstand der Erfindung ist auch ein Verfahren zur Isolierung einer entsprechenden Zellpopulation mit folgenden Schritten:

entweder

- Entnahme einer Probe aus dem Hirn
- Isolieren der neuralen Vorläuferzellen unter Ver- 20 wendung der angegebenen Marker

oder

- Differenzierung von embryonalen Stammzellen zu neuralen Vorläuferzellen,
- Isolieren der neuralen Vorläuferzellen unter Verwendung der angegebenen Marker

oder

- Trans-Differenzierung von adulten, nicht neuralen Stammzellen zu neuralen Vorläuferzellen.
- Isolieren der neuralen Vorläuferzellen unter Verwendung der angegebenen Marker

oder

- Differenzierung von adulten, neuralen Stammzellen zu neuralen Vorläuferzellen,
- Isolieren der neuralen Vorläuferzellen unter Verwendung der angegebenen Marker

oder

- Differenzierung von immortalisierten Zellen zu neuralen Vorläuferzellen,
- Isolieren der neuralen Vorläuferzellen unter Verwendung der angegebenen Marker.

[0019] "Unter Verwendung der angegebenen Marker" bedeutet, dass die Zellen isoliert werden, die positiv für mindestens einen der Marker aus der Liste A und B sind, wobei mehrere positive Marker und die Abwesenheit von Markern der Liste C bevorzugt werden. Die Isolierung kann beispielsweise durch FACS Analyse erfolierung kann beispielsweise durch Erfolierung kann beispielsweise durch Erfolierung kann beispielswe

gen. Die durch die Verfahren erhältlichen Zellen sind ebenfalls Gegenstand der Erfindung.

[0020] Ein weiterer Gegenstand der Erfindung ist die Verwendung mindestens eines Markers ausgewählt aus der Liste A oder Liste B zu Identifizierung oder Isolierung von neuralen Vorläuferzellen.

[0021] Gegenstand ist weiterhin ein Antikörper gegen einen Marker aus der Liste A, B oder C, ein Diagnostikmittel enthaltend mindestens einen, bevorzugt zwei oder mehr Substanzen zur Erkennung der Marker der Liste A, B oder C sowie ein Arzneimittel enthaltend die erfindungsgemäße Zellpopulation.

[0022] Solche Arzneimittel könnten wie oben dargestellt zur Behandlung von neurologischen Krankheiten wie Alzheimer, Parkinson, Folgen von Schädelhirntraumata oder Schlaganfall eingesetzt werden.

[0023] Ein weiterer Gegenstand ist eine Zellpopulation, bei der mindestens 5% der Zellen neurale Stammzellen sind, die wenigstens einen der in Liste D oder Liste E aufgeführten Marker aufweisen.

[0024] Vorzugsweise weisen entsprechende neurale Stammzellen mindestens zwei, bevorzugt mindestens drei, mindestens vier und noch mehr bevorzugt mindestens fünf der in Liste D oder Liste E aufgeführten Marker auf.

[0025] In besonders bevorzugten Ausführungsformen weisen entsprechende neurale Stammzellen keinen der in Liste A oder Liste C aufgeführten Marker auf. [0026] Der Gehalt an neuralen Stammzellen in der Zellpopulation ist möglichst hoch, bevorzugt mindestes 10%, mehr bevorzugt mindestes 25%, mindestens 50%, und am meisten bevorzugt mindestens 90%.

[0027] Entsprechende Zellpopulation sind aus Hirngewebe erhältlich. In einer Ausführungsform handelt es sich um eine murine Zellpopulation.

[0028] Gegenstand ist weiterhin ein Verfahren zur Isolierung der Zellpopulation. Dies ist erhältlich entweder durch

- Entnahme einer Probe aus dem Hirn
 - Isolieren der neuralen Stammzellen unter Verwendung der angegebenen Marker

45 oder

- Differenzierung von embryonalen Stammzellen zu neuralen Stammzellen,
- Isolieren der neuralen Stammzellen unter Verwendung der angegebenen Marker

oder

- Trans-Differenzierung von adulten, nicht neuralen Stammzellen zu neuralen Stammzellen.
 - Isolieren der neuralen Stammzellen unter Verwen-

dung der angegebenen Marker

oder

- De-Differenzierung von adulten, neuralen Vorläuferzellen zu neuralen Stammzellen,
- Isolieren der neuralen Stammzellen unter Verwendung der angegebenen Marker

oder

- Differenzierung von immortalisierten Zellen zu neuralen Stammzellen,
- Isolieren der neuralen Stammzellen unter Verwendung der angegebenen Marker.

[0029] Die Isolierung erfolgt wie oben bei den neuralen Vorläuferzellen angegeben. Auch die auf diesem Wege erhältlichen neuralen Stammzellen sind Gegenstand der Erfindung.

[0030] Gegenstand der Erfindung ist weiterhin ein Antikörper gegen einen Marker aus der Liste D, E, ein Diagnostikmittel enthaltend mindestens einen, bevorzugt zwei oder mehr Substanzen zur Erkennung der Marker der Liste D, E, A oder C sowie ein Arzneimittel enthaltend die erfindungsgemäße Zellpopulation.

[0031] Solche Arzneimitteln können wie dargestellt zur Behandlung von neuronalen Krankheiten wie Alzheimer, Parkinson, Folgen von Schädelhirntraumata oder Schlaganfall eingesetzt werden.

Beispiele

A. Isolierung von embryonaler Stammzellen

[0032] Murine embryonale Stammzellen proliferieren klonal in vitro und sind aus diesem Grunde in großer Menge und hochreiner Form isolierbar. Nach dem Stand der Technik werden diese in Anwesenheit von LIF auf primären embryonalen Fibroblasten gehalten und regelmäßig durch die Generierung von hochgradig keimbahnkompetenten chimären Mäusen auf ihre Qualität überprüft. Unter normalen Kulturbedingungen beträgt das Verhältnis ES-Zellen zu kontaminierenden Fibroblasten etwa 200:1. Um auch diese minoritäre Komponente zu eliminieren, wurden die ES-Zellen vor der RNA-Päparation für zwei Passagen (vier Tage) auf gelatinisierten Kulturplatten bei erhöhter LIF-Konzentration gehalten. Dies ermöglicht eine Reduktion der kontaminierenden Fibroblasten auf etwa 0,01% der Gesamtpopulation.

B. Isolierung von neuronalen Vorläuferzellen aus dem adulten Mausgehirn.

[0033] In der subventrikulären Zone des adulten Vor-

derhirns von Vertebraten werden permanent große Mengen von neuralen Vorläuferzellen gebildet (wahrscheinlich < 50000 Zellen/ Tag). Diese Zellen benutzen einen präzise definierten Migrationsweg und eine spezielle Form der Translokation (*Chain migration*) um in den Bulbus olfaktorius zu gelangen. Im Bulbus olfaktorius angelangt differenzieren diese Vorläuferzellen normalerweise in inhibitorische (GABA-erge) Interneurone. Unter bestimmten experimentellen Bedingungen wurde ihre Differenzierung in Oligodendrozyten und Astrozy-

6

ten gezeigt.

[0034] Neurale Vorläufer, die einen Differenzierungszustand zwischen einer neuralen Stammzelle und einem terminal differenzierten Neuron repräsentieren, exprimieren spezifisch eine Form des neuralen Zelladhäsionsmoleküls NCAM, die eine spezielle post-translationelle Modifikation aufweist. Diese Modifikation besteht aus der Glykosylierung des Proteins mit a-2,8 verknüpfter Polysialylsäure (PSA). Ein spezifischer Antikörper gegen dieses Glykoepitop (Chazal et al., 2000) erlaubte die hochreine Isolierung der Zielpopulation aus dissozierten Vorderhirngewebe durch FACS (Fluorescence

5 C. Molekulargenetische Analyse

Activated Cell Sorting).

[0035] Embryonale Stammzellen und neuronale Vorläuferzellen wurden in einem genomweiten Screen mit der Methode SAGE (Serial Analysis of Gene Expression) analysiert.

[0036] Die Genexpressionsprofile der beiden Zell-Populationen wurden unter Anwendung bioinformatischer Verfahrensweisen mit Maus-Hirn-SAGE-Datenbanken verglichen, um molekulare Marker zu identifizieren, die charakteristisch für embryonale Stammzellen und neuronale Vorläuferzellen sind.

[0037] Mit Hilfe der Microarray technologie wurde die Expression der Gene bestätigt.

[0038] Durch in situ-Hybridisierung in Maushirn und an embryonalen Stammzellen wurde die zelluläre Lokalisation einiger der identifizierten Gene bestimmt. Diese Ergebnisse belegen, dass spezifische Markergene identifiziert werden konnten.

5 Liste A: Positivmarker neurale Vorläuferzellen (1.) und Negativmarker 2 neurale Stammzellen;

ES-Zellen -; PSA-NCAM +; Adult brain -

0 [0039]

	Mm.8884	nuclear factor of kappa light chain gene
		enhancer in B-cells inhibitor, alpha
	Mm.8180	lymphocyte antigen 6 complex, locus A
5	Mm.6238	SRY-box containing gene 11
	Mm.517	(Manual) Manic fringe protein, putative
		secreted glycosyltransferase, notch
		modulator

7

Mm.4919 Mm.4727	DNA segment, human D4S114 seizure related gene 6		Liste B: Posi	itivmarker neurale Vorläuferzellen (2.);	
Mm.45769	ESTs		ES-Zellen -/+; PSA-NCAM +; Adult brain -		
Mm.44490 Mm.42948	RIKEN cDNA 6330415M09 gene peroxiredoxin 2	5	[0040]		
Mm.4022 Mm.3940	RIKEN cDNA 1110033C18 gene lethal giant larvae homolog		Mm.911	high mobility group puolessemat hin	
Mm.37835	ribosomal protein L7		WIIII.911	high mobility group nucleosomal bin- ding domain 2	
Mm.3779	RIKEN cDNA 2300006C11 gene		Mm.89136	H3 histone, family 3A	
Mm.340	high mobility group box 3	10	Mm.741	fatty acid binding protein 5, epidermal	
Mm.32902	ESTs, Weakly similar to S26689 hypo-		Mm.7286	C-terminal binding protein 1	
	thetical protein hc1 - mouse		Mm.7141	proliferating cell nuclear antigen	
Mm.3268	ubiquitin-conjugating enzyme E2I		Mm.6840	RIKEN cDNA 5730507C05 gene	
Mm.31436	myeloid ecotropic viral integration site-		Mm.6787	splicing factor, arginine/serine-rich 3	
•	related gene 1	15		(SRp20)	
Mm.297	actin, beta, cytoplasmic		Mm.6417	CD24a antigen	
Mm.29558	expressed sequence Al426163		Mm.6343	nucleophosmin 1	
Mm.29014	T-cell lymphoma invasion and metasta-		Mm.482	Jun oncogene	
	sis 2		Mm.43871	expressed sequence AW046487	
Mm.28842	chloride channel 3	20	Mm.43213	RIKEN cDNA 9030402K04 gene	
Mm.28824	Mus musculus, clone IMAGE:4504748,		Mm.42767	ribosomal protein S17	
	mRNA		Mm.4269	transcription factor 4	
Mm.28275	RNA binding motif protein, X chromoso-		Mm.40715	RIKEN cDNA 1110038H03 gene	
	me		Mm.40715	RIKEN cDNA 1110038H03 gene	
Mm.28149	RIKEN cDNA 3110003A17 gene	25	Mm.4071	laminin receptor 1 (67kD, ribosomal pro-	
Mm.28148	chromobox homolog 3 (Drosophila HP1			tein SA)	
M 07040	gamma)		Mm.4025	nuclear factor I/B	
Mm.27816	hexosaminidase B		Mm.372	ribosomal protein S26	
Mm.2769	MARCKS-like protein		Mm.3487	ribosomal protein L30	
Mm.22171	calponin 3, acidic	30	Mm.3381	ribosomal protein S8	
Mm.220923	RIKEN cDNA 6530406007 gene		Mm.31051	RIKEN cDNA 2610003J05 gene	
Mm.21740	heterogeneous nuclear ribonucleopro-		Mm.30120	ribosomal protein S27-like	
	tein H1		Mm.30011	ribosomal protein S23	
Mm.206085	expressed sequence Al854782		Mm.29911	RIKEN cDNA 3200001M24 gene	
Mm.205996	EST AA087124	35	Mm.2966	isocitrate dehydrogenase 2 (NADP+),	
Mm.200858	RIKEN cDNA 2410129E14 gene			mitochondrial	
Mm.199500	expressed sequence Al844617		Mm.29580	superiorcervical ganglia, neural specific	
Mm.195901	ribosomal protein L35a			10	
Mm.194965	EST		Mm.2958	expressed sequence Al843786	
Mm.19101	DEAD (aspartate-glutamate-alanine-	40	Mm.28985	ribosomal protein L27	
Mm.19016	aspartate) box polypeptide 5		Mm.28869	ESTs	
	drebrin 1		Mm.27927	heterogeneous nuclear ribonucleopro-	
Mm.18789	SRY-box containing gene 4			tein A1	
Mm.186740	ESTs		Mm.27669	small nuclear ribonucleoprotein E	
Mm.18516	H3 histone, family 3B	45	Mm.2756.	high mobility group nucleosomal bin-	
Mm.181959	early growth response 1			ding domain 1	
Mm.181847	prefoldin 5		Mm.27141	Rac GTPase-activating protein 1	
Mm. 16421	high mobility group box 1		Mm.2591	RNA binding motif protein 3	
Mm.15534	interleukin 1 alpha		Mm.24083	Mus musculus, Similar to TAR DNA bin-	
Mm.13725	Paneth cell enhanced expression	50		ding protein, clone MGC: 19284	
Mm.12871	doublecortin		N 040000	IMAGE:4016437, mRNA, complete cds	
Mm.127662	ESTS Mus museulus Similar ta DIKEN aDNA		Mm.219668	RIKEN cDNA 2610209F03 gene	
Mm.12412	Mus musculus, Similar to RIKEN cDNA 2810407E23 gene, clone IMAGE:		Mm.21841	splicing factor, arginine/serine-rich 2 (SC-35)	
	4489006, mRNA, partial cds	55	Mm.218240	Mus musculus, clone IMAGE:5342828,	
			Mm.21740	mRNA, partial cds heterogeneous nuclear ribonucleopro-	
				tein H1	

9	EP 1 529 838 A1	10

Mm.213020	(Manual) 60S ribosomal protein L32 (RPL32)		Mm.6660 Mm.6586	small inducible cytokine A27 Mus musculus, clone MGC:6299 IMAGE:
Mm.2115	heterogeneous nuclear ribonucleopro-		14 0505	2654341, mRNA, complete cds
Mm.196611	tein U synapsin I	5	Mm.6565 Mm.65337	FK506 binding protein 8 (38 kDa)
Mm.19187	prothymosin alpha	,	WIII1.03337	Mus musculus, clone MGC:28924 IMAGE:3481738, mRNA, complete cds
Mm.18789	SRY-box containing gene 4		Mm.648	prion protein
Mm.186499	_ ·		Mm.638	ESTs
	superfamily containing leucinerich re-		Mm.544	phosphoprotein enriched in astrocytes 15
	peat	10	Mm.5264	ESTs, Highly similar to FEZ1_RAT FA-
Mm.18516	H3 histone, family 3B			SCICULATION AND ELONGATION
Mm.180873	RIKEN cDNA 2510019J09 gene			PROTEIN ZETA 1 (ZYGIN I)
Mm.1775	hematological and neurological expres-		Mm.5259	(Manual assignment) probably myelin-
	sed sequence 1			associated oligodendrocyte basic protein
Mm.1703	tubulin, beta 5	15		MOBP
Mm.16775	ribosomal protein S24		Mm.5249	copine 6
Mm.16767	heterogeneous nuclear ribonucleopro-		Mm.52	RIKEN cDNA 1810033A19 gene
M 40500	tein A2/B1		Mm.5195	complexin 1
Mm.16596	B-cell translocation gene 1, anti-prolife-		Mm.5153	neurotensin receptor 2
M + 40070	rative	20	Mm.5023	Purkinje cell protein 4
Mm.148973 Mm.142872	RIKEN cDNA 3010025E17 gene		Mm.4923	ESTs
WIIII. 142072	heterogeneous nuclear ribonucleopro- tein K		Mm.4921	glutamate receptor, ionotropic, AMPA2
Mm.142729	thymosin, beta 4, X chromosome		Mm.4920	(alpha 2)
Mm.140380	ribosomal protein L23	25	WIIII.4920	glutamate receptor, ionotropic, AMPA1 (alpha 1)
Mm.140	protein phosphatase 1, regulatory (inhi-		Mm.4870	synaptosomal-associated protein, 91
	bitor) subunit 14B		1411114070	kDa
Mm. 12858	eukaryotic translation initiation factor		Mm.4857	calcium/calmodulin-dependent protein
	4A1			kinase II, beta
		30	Mm.4762	kinesin heavy chain member 1A
	gativmarker 1 neurale Stammzellen und		Mm.4705	(Manual) probably in far 3'-UTR of com-
Negativmar	ker neurale Vorläuferzellen;			plexin-2 cDNA
			Mm.46764	RIKEN cDNA 4833409J18 gene
ES-Zellen -;	PSA-NCAM -; Adult brain +		Mm.4657	amyloid beta (A4) precursor protein-bin-
[0044]		35		ding, family A, member 2
[0041]			Mm.4651	kinesin-associated protein 3
Mm.98	proteasome (prosome, macropain) subu-		Mm.45951	RIKEN cDNA 1200016B17 gene
141111.30	nit, beta type 6		Mm.4550	ATPase, Na+/K+ transporting, beta 1 po- lypeptide
Mm.9745	lactate dehydrogenase 2, B chain	40	Mm.4550	ATPase, Na+/K+ transporting, beta 1 po-
Mm.970	creatine kinase, mitochondrial 1, ubiqui-		1411111000	ATT 430, NATATE ITALISPOILING, DETA 1 PO-
				lypeptide
Mm.891	tous		Mm.4537	lypeptide NADH dehydrogenase (ubiquinone) 1
141111111111111111111111111111111111111	tous kinesin family member C2		Mm.4537	NADH dehydrogenase (ubiquinone) 1
Mm.88833			Mm.4537 Mm.44355	•••
	kinesin family member C2	45		NADH dehydrogenase (ubiquinone) 1 beta subcomplex, 9
	kinesin family member C2 Mus musculus strain ILS K-Cl cotrans- porter (Slc12a5) mRNA, complete cds BM88 antigen	45	Mm.44355	NADH dehydrogenase (ubiquinone) 1 beta subcomplex, 9 RIKEN cDNA 6430514L14 gene
Mm.88833 Mm.87027 Mm.8688	kinesin family member C2 Mus musculus strain ILS K-Cl cotrans- porter (Slc12a5) mRNA, complete cds BM88 antigen RIKEN cDNA 0610011B04 gene	45	Mm.44355	NADH dehydrogenase (ubiquinone) 1 beta subcomplex, 9 RIKEN cDNA 6430514L14 gene synaptosomal-associated protein, 25
Mm.88833 Mm.87027 Mm.8688 Mm.86654	kinesin family member C2 Mus musculus strain ILS K-Cl cotransporter (Slc12a5) mRNA, complete cds BM88 antigen RIKEN cDNA 0610011B04 gene microtubule-associated protein 6	45	Mm.44355 Mm.4435	NADH dehydrogenase (ubiquinone) 1 beta subcomplex, 9 RIKEN cDNA 6430514L14 gene synaptosomal-associated protein, 25 kDa
Mm.88833 Mm.87027 Mm.8688 Mm.86654 Mm.848	kinesin family member C2 Mus musculus strain ILS K-Cl cotransporter (Slc12a5) mRNA, complete cds BM88 antigen RIKEN cDNA 0610011B04 gene microtubule-associated protein 6 testis expressed gene 261		Mm.44355 Mm.4435 Mm.44244	NADH dehydrogenase (ubiquinone) 1 beta subcomplex, 9 RIKEN cDNA 6430514L14 gene synaptosomal-associated protein, 25 kDa open reading frame 12 ESTs Mus musculus, ATPase, Na+K+ trans-
Mm.88833 Mm.87027 Mm.8688 Mm.86654 Mm.848 Mm.806	kinesin family member C2 Mus musculus strain ILS K-Cl cotransporter (Slc12a5) mRNA, complete cds BM88 antigen RIKEN cDNA 0610011B04 gene microtubule-associated protein 6 testis expressed gene 261 CD 81 antigen	45 50	Mm.44355 Mm.4435 Mm.44244 Mm.44107	NADH dehydrogenase (ubiquinone) 1 beta subcomplex, 9 RIKEN cDNA 6430514L14 gene synaptosomal-associated protein, 25 kDa open reading frame 12 ESTs Mus musculus, ATPase, Na+K+ transporting, alpha 3 subunit, clone MGC:
Mm.88833 Mm.87027 Mm.8688 Mm.86654 Mm.848	kinesin family member C2 Mus musculus strain ILS K-Cl cotransporter (Slc12a5) mRNA, complete cds BM88 antigen RIKEN cDNA 0610011B04 gene microtubule-associated protein 6 testis expressed gene 261 CD 81 antigen ESTs, Weakly similar to simple repeat se-		Mm.44355 Mm.4435 Mm.44244 Mm.44107	NADH dehydrogenase (ubiquinone) 1 beta subcomplex, 9 RIKEN cDNA 6430514L14 gene synaptosomal-associated protein, 25 kDa open reading frame 12 ESTs Mus musculus, ATPase, Na+K+ transporting, alpha 3 subunit, clone MGC: 27631 IMAGE:4506376, mRNA, comple-
Mm.87027 Mm.8688 Mm.86654 Mm.848 Mm.806 Mm.80123	kinesin family member C2 Mus musculus strain ILS K-Cl cotransporter (Slc12a5) mRNA, complete cds BM88 antigen RIKEN cDNA 0610011B04 gene microtubule-associated protein 6 testis expressed gene 261 CD 81 antigen ESTs, Weakly similar to simple repeat sequence-containing transcript		Mm.44355 Mm.4435 Mm.44244 Mm.44107 Mm.44101	NADH dehydrogenase (ubiquinone) 1 beta subcomplex, 9 RIKEN cDNA 6430514L14 gene synaptosomal-associated protein, 25 kDa open reading frame 12 ESTs Mus musculus, ATPase, Na+K+ transporting, alpha 3 subunit, clone MGC: 27631 IMAGE:4506376, mRNA, complete cds
Mm.88833 Mm.87027 Mm.8688 Mm.86654 Mm.848 Mm.806 Mm.80123	kinesin family member C2 Mus musculus strain ILS K-Cl cotransporter (Slc12a5) mRNA, complete cds BM88 antigen RIKEN cDNA 0610011B04 gene microtubule-associated protein 6 testis expressed gene 261 CD 81 antigen ESTs, Weakly similar to simple repeat sequence-containing transcript aldolase 3, C isoform		Mm.44355 Mm.4435 Mm.44244 Mm.44107 Mm.44101	NADH dehydrogenase (ubiquinone) 1 beta subcomplex, 9 RIKEN cDNA 6430514L14 gene synaptosomal-associated protein, 25 kDa open reading frame 12 ESTs Mus musculus, ATPase, Na+K+ transporting, alpha 3 subunit, clone MGC: 27631 IMAGE:4506376, mRNA, complete cds myc box dependent interacting protein 1
Mm.87027 Mm.8688 Mm.86654 Mm.848 Mm.806 Mm.80123	kinesin family member C2 Mus musculus strain ILS K-Cl cotransporter (Slc12a5) mRNA, complete cds BM88 antigen RIKEN cDNA 0610011B04 gene microtubule-associated protein 6 testis expressed gene 261 CD 81 antigen ESTs, Weakly similar to simple repeat sequence-containing transcript aldolase 3, C isoform tubulin, beta 4		Mm.44355 Mm.44244 Mm.44107 Mm.44101 Mm.4383 Mm.43786	NADH dehydrogenase (ubiquinone) 1 beta subcomplex, 9 RIKEN cDNA 6430514L14 gene synaptosomal-associated protein, 25 kDa open reading frame 12 ESTs Mus musculus, ATPase, Na+K+ transporting, alpha 3 subunit, clone MGC: 27631 IMAGE:4506376, mRNA, complete cds myc box dependent interacting protein 1 cytochrome c oxidase, subunit VIIc
Mm.88833 Mm.87027 Mm.8688 Mm.86654 Mm.848 Mm.806 Mm.80123 Mm.7729 Mm.7420	kinesin family member C2 Mus musculus strain ILS K-Cl cotransporter (Slc12a5) mRNA, complete cds BM88 antigen RIKEN cDNA 0610011B04 gene microtubule-associated protein 6 testis expressed gene 261 CD 81 antigen ESTs, Weakly similar to simple repeat sequence-containing transcript aldolase 3, C isoform	50	Mm.44355 Mm.44244 Mm.44107 Mm.44101 Mm.4383 Mm.43786 Mm.43749	NADH dehydrogenase (ubiquinone) 1 beta subcomplex, 9 RIKEN cDNA 6430514L14 gene synaptosomal-associated protein, 25 kDa open reading frame 12 ESTs Mus musculus, ATPase, Na+K+ transporting, alpha 3 subunit, clone MGC: 27631 IMAGE:4506376, mRNA, complete cds myc box dependent interacting protein 1 cytochrome c oxidase, subunit VIIc RIKEN cDNA 3100001N19 gene
Mm.88833 Mm.87027 Mm.8688 Mm.86654 Mm.848 Mm.806 Mm.80123 Mm.7729 Mm.7420 Mm.7363	kinesin family member C2 Mus musculus strain ILS K-CI cotransporter (Slc12a5) mRNA, complete cds BM88 antigen RIKEN cDNA 0610011B04 gene microtubule-associated protein 6 testis expressed gene 261 CD 81 antigen ESTs, Weakly similar to simple repeat sequence-containing transcript aldolase 3, C isoform tubulin, beta 4 beta-spectrin 3	50	Mm.44355 Mm.44244 Mm.44107 Mm.44101 Mm.4383 Mm.43786	NADH dehydrogenase (ubiquinone) 1 beta subcomplex, 9 RIKEN cDNA 6430514L14 gene synaptosomal-associated protein, 25 kDa open reading frame 12 ESTs Mus musculus, ATPase, Na+K+ transporting, alpha 3 subunit, clone MGC: 27631 IMAGE:4506376, mRNA, complete cds myc box dependent interacting protein 1 cytochrome c oxidase, subunit VIIc
Mm.88833 Mm.87027 Mm.8688 Mm.86654 Mm.848 Mm.806 Mm.80123 Mm.7729 Mm.7420 Mm.7363 Mm.726	kinesin family member C2 Mus musculus strain ILS K-CI cotransporter (Slc12a5) mRNA, complete cds BM88 antigen RIKEN cDNA 0610011B04 gene microtubule-associated protein 6 testis expressed gene 261 CD 81 antigen ESTs, Weakly similar to simple repeat sequence-containing transcript aldolase 3, C isoform tubulin, beta 4 beta-spectrin 3 basigin	50	Mm.44355 Mm.44244 Mm.44107 Mm.44101 Mm.4383 Mm.43786 Mm.43749 Mm.43721	NADH dehydrogenase (ubiquinone) 1 beta subcomplex, 9 RIKEN cDNA 6430514L14 gene synaptosomal-associated protein, 25 kDa open reading frame 12 ESTs Mus musculus, ATPase, Na+K+ transporting, alpha 3 subunit, clone MGC: 27631 IMAGE:4506376, mRNA, complete cds myc box dependent interacting protein 1 cytochrome c oxidase, subunit VIIc RIKEN cDNA 3100001N19 gene small nuclear ribonucleoprotein N

	EP	1	529	838	A1	
--	----	---	-----	-----	-----------	--

	lypeptide 1		Mm.3974	ubiquitin specific protease 4 (proto-onco-
Mm.4339	laminin, alpha 5			gene)
Mm.43330	RIKEN cDNA 0610025G13 gene		Mm.39548	expressed sequence AI839779
Mm.43278	olfactomedin 1		Mm.3951	thymus cell antigen 1, theta
Mm.43278	olfactomedin 1	5	Mm.3915	myelin-associated oligodendrocytic ba-
Mm.4296	synovial sarcoma translocation, Chromo-			sic protein
	some 18		Mm.39040	myelin and lymphocyte protein, T-cell dif-
Mm.42949	RIKEN cDNA 1110012005 gene			ferentiation protein
Mm.42948	peroxiredoxin 2		Mm.38994	RIKEN cDNA 2600001N01 gene
Mm.42829	selenoprotein W, muscle 1	10	Mm.38993	calsyntenin 1
Mm.4266	integral membrane protein 2B		Mm.38551	calcium binding protein 1
Mm.4266	integral membrane protein 2B		Mm.38469	amyloid beta (A4) precursor protein-bin-
Mm.4263	cystatin C			ding, family B, member 1
Mm.425	histidine triad nucleotide binding protein		Mm.38438	RIKEN cDNA 1200009K17 gene
Mm.42255	ATPase, Ca++ transporting, cardiac	15	Mm.38421	(Manual assignment) ATPase, Na+K+
	muscle, slow twitch 2			transporting, alpha polypeptide
Mm.41926	NADH dehydrogenase (ubiquinone) 1 al-		Mm.38421	(Manual assignment) ATPase, Na+K+
	pha subcomplex, 4			transporting, alpha polypeptide
Mm.41925	RIKEN cDNA 1810034B16 gene		Mm.3840	flotillin 2
Mm.41918	RIKEN cDNA 1110063G11 gene	20	Mm.38248	sialyltransferase 9 (CMP-NeuAc:lacto-
Mm.41911	cytochrome P450, 46 (cholesterol 24-hy-			sylceramide alpha-2,3-sialyltransferase)
	droxylase)		Mm.38036	ESTs, Moderately similar to
Mm.41893	RIKEN cDNA 6330408G06 gene			NX1A_MOUSE_2
Mm.41791	glycoprotein m6b		Mm.38036	ESTs, Moderately similar to
Mm.41752	expressed sequence Al847934	25		NX1A_MOUSE_2
Mm.41735	RIKEN cDNA 2300004C15 gene		Mm.37462	ESTs, Weakly similar to CA11 RAT COL-
Mm.41719	RIKEN cDNA 2610507A21 gene			LAGEN ALPHA 1(I) CHAIN
Mm.41711	Mus musculus, clone IMAGE:3499845,		Mm.37214	transferrin
	mRNA, partial cds		Mm.36275	DNA segment, Chr 11, Brigham & Wo-
Mm.41694	ESTs	30		men's Genetics 0517 expressed
Mm.41692	ESTs, Weakly similar to F59F4.2.p		Mm.3624	guanylate kinase 1
Mm.41642	regulator of G-protein signaling 4		Mm.35837	RIKEN cDNA 2510006D16 gene
Mm.41630	RIKEN cDNA 0710001E10 gene		Mm.35837	RIKEN cDNA 2510006D16 gene
Mm.41604	ESTs, Weakly similar to VAV3_MOUSE		Mm.3544	calcium channel, voltage-dependent, be-
	VAV-3 PROTEIN	35		ta 3 subunit
Mm.41603	expressed sequence Al891706		Mm.35439	secreted acidic cysteine rich glycoprotein
Mm.41603	expressed sequence Al891706		Mm.35270	Ly6/neurotoxin 1
Mm.41602	RIKEN cDNA 3110050007 gene		Mm.3479	ATPase, H+ transporting, lysosomal
Mm.41602	RIKEN cDNA 3110050007 gene			21kDa, V0 subunit B
Mm.4137	chromogranin A	40	Mm.34695	actin related protein 2/3 complex, subunit
Mm.41354	ESTs			1A (41 kDa)
Mm.41277	RIKEN cDNA 1110020M21 gene		Mm.34246	calmodulin 1
Mm.41248	ESTs		Mm.3363	prosaposin
Mm.41190	RIKEN cDNA 1700112L09 gene		Mm.3360	tyrosine 3-monooxygenase/tryptophan
Mm.40863	expressed sequence AW049870	45		5-monooxygenase activation protein, ze-
Mm.40738	RIKEN cDNA 2900072M03 gene			ta polypeptide
Mm.40621	ESTs, Moderately similar to		Mm.33117	ESTs
	Y552_HUMAN HYPOTHETICAL PRO-		Mm.3308	tyrosine 3-monooxygenase/tryptophan
	TEIN KIAA0552			5-monooxygenase activation protein, eta
Mm.40472	expressed sequence Al835002	50		polypeptide
Mm.40443	RIKEN cDNA 4930488B01 gene		Mm.3292	glutamate receptor, ionotropic, NMDA1
Mm.40124	phosphodiesterase 10A			(zeta 1)
Mm.40059	ESTs, Weakly similar to SP62 MOUSE		Mm.3229	ribosomal protein L26
	SPLICEOSOME ASSOCIATED PROTE-		Mm.32191	gamma-aminobutyric acid (GABA-B) re-
	IN 62	5 5		ceptor, 1
Mm.39857	RIKEN cDNA 2900074L19 gene		Mm.31395	carboxypeptidase E
Mm.39803	expressed sequence Al841080		Mm.3123	comichon-like (Drosophila)
Mm.39752	RIKEN cDNA 2900041A09 gene		Mm.31025	RIKEN cDNA 2310015K15 gene

EP 1	529	838	A1

Mm.30412 Mm.30355	RIKEN cDNA 5430400P17 gene (Manual) KIF5A Neuronal Kinesin heavy chain		Mm.29230 Mm.29227 Mm.29205	RIKEN cDNA 1500017E18 gene RIKEN cDNA 2300002D11 gene bruno-like 4, RNA binding protein (Droso-
Mm.30266	hemoglobin, beta adult major chain			phila)
Mm.30266	hemoglobin, beta adult major chain	5	Mm.29205	bruno-like 4, RNA binding protein (Droso-
Mm.30206	ATPase, H+ transporting, lysosomal			phila)
	34kD, V1 subunit D		Mm.2918	megakaryocyte-associated tyrosine ki-
Mm.30156	protease, serine, 11 (Igf binding)			nase
Mm.30155	ATPase, H+ transporting, lysosomal		Mm.29141	RIKEN cDNA 0710008N11 gene
Mm.30150	16kD, V0 subunit C	10	Mm.29124	phosphatidic acid phosphatase type 2B
Mm.30126	RIKEN cDNA 1010001M12 gene		Mm.29075	(Manual) Reticulon 1 protein, major inter-
Mm.30085	membrane interacting protein of RGS16		N= 00007	nal tag
141111.30003	aldo-keto reductase family 1, member A4 (aldehyde reductase)		Mm.29027 Mm.29027	SPARC-like 1 (mast9, hevin)
Mm.30072	cytochrome c oxidase subunit VIIa poly-	15	Mm.29027	SPARC-like 1 (mast9, hevin)
WIIII.00072	peptide 2-like		WIII1.2902	protein tyrosine phosphatase, receptor- type, N
Mm.30059	myristoylated alanine rich protein kinase		Mm.28955	RIKEN cDNA 4930570C03 gene
	C substrate		Mm.28650	RAB6, member RAS oncogene family
Mm.29976	septin 5		Mm.28650	RAB6, member RAS oncogene family
Mm.29965	RIKEN cDNA 2410104119 gene	20	Mm.28643	vesicle-associated membrane protein 2
Mm.29947	serine/threonine kinase 11		Mm.28561	protein kinase C, zeta
Mm.29939	RIKEN cDNA 1010001N11 gene		Mm.28518	type I transmembrane protein Fn14
Mm.29937	(Manual assignment) polymorphism of		Mm.28357	microtubule-associated protein 1 light
	Mm.29937 ESTs, Weakly similar to pre-			chain 3
	dicted using Genefinder	25	Mm.2815	RIKEN cDNA 1110021H02 gene
Mm.29921	RAS protein-specific guanine nucleotide-		Mm.28107	ectonucleotide pyrophosphatase/phos-
	releasing factor 1			phodiesterase 2
Mm.2992	(Manual assignment) MBP myelin basic		Mm.28058	NADH dehydrogenase (ubiquinone) 1
	protein			beta subcomplex 5
Mm.29870	integral membrane protein 3	30	Mm.27886	RIKEN cDNA 2410011G03 gene
Mm.29867	NADH dehydrogenase (ubiquinone) 1 al-			3.7.1
	pha subcomplex 2		Mm.27608	Mus musculus, Similar to chromosome
Mm.29857	(Manual) Neurogranin			9 open reading frame 16, clone MGC:
Mm.29852	Mus musculus, clone IMAGE:5102170,			19388 IMAGE:2812475, mRNA, com-
	mRNA, partial cds	35		plete cds
Mm.29846	Mus musculus, Similar to NDRG family,		Mm.2755	calbindin 2
	member 4, clone MGC:7067 IMAGE:		Mm.27499	RIKEN cDNA 2010004E11 gene
	3156802, mRNA, complete cds		Mm.27407	RecQ protein-like
Mm.29842	NADH dehydrogenase flavoprotein 1		Mm.27256	discs, large homolog 4 (Drosophila)
Mm.29823	microsomal glutathione S-transferase 3	40	Mm.2720	mitogen activated protein kinase 8 in-
Mm.29807	ubiquitin carboxy-terminal hydrolase L1			teracting protein
Mm.29807	ubiquitin carboxy-terminal hydrolase L1		Mm.27114	RIKEN cDNA 0610043B10 gene
Mm.29771	ATPase, H+ transporting, lysosomal		Mm.27087	RIKEN cDNA 2010012C24 gene
Mm 20717	70kD, V1 subunit A, isoform 1	45	Mm.27005	visinin-like 1
Mm.29717	3-monooxgenase/tryptophan 5-monoox- genase activation protein, gamma poly-	45	Mm.26633	PH domain containing protein in retina 1
	peptide		Mm.26633	PH domain containing protein in retina 1
Mm.29711	adrenergic receptor kinase, beta 1		Mm.26550 Mm.2645	phosphofructokinase, muscle
Mm.297	actin, beta, cytoplasmic		WIII1.2043	eukaryotic translation elongation factor 1 alpha 2
Mm.29633	RIKEN cDNA 1810008021 gene	50	Mm.2635	pyruvate kinase 3
Mm.29600	Mus musculus, clone IMAGE:3964267.		Mm.2619	cholecystokinin
	mRNA		Mm.25849	RIKEN cDNA 2010003014 gene
Mm.2948	H2-K region expressed gene 2		Mm.25738	RIKEN cDNA 2900002P20 gene
Mm.29477	SCAN domain-containing 1		Mm.25228	ring finger protein 11
Mm.29415	RIKEN cDNA 1810011001 gene	55	Mm.25203	NCK-associated protein 1
Mm.29362	expressed sequence Al414999		Mm.2496	internexin neuronal intermediate fila-
Mm.29344	tumor differentially expressed 1, like			ment protein, alpha
Mm.29330	expressed sequence AI853543		Mm.24482	RIKEN cDNA 5730460C18 gene
				Ç

ΕP	1	529	838	A1	

Mm.2446	synaptotagmin 4			pha o
Mm.24376	Mus musculus mRNA for calsyntenin-3		Mm.20964	guanine nucleotide binding protein, al-
	(Cs3 gene)			pha o
Mm.2411	Ras-GTPase-activating protein (GAP		Mm.2082	apolipoprotein D
	<120>) SH3-domain binding protein 2	5	Mm.206218	Mus musculus, Similar to hypothetical
Mm.24092	N-ethylmaleimide sensitive fusion pro-			protein FLJ22237, clone MGC:27683
	tein			IMAGE:4913322, mRNA, complete cds
Mm.24092	N-ethylmaleimide sensitive fusion pro-		Mm.2060	RIKEN cDNA 2900010105 gene
	tein		Mm.20472	vertebrate homolog of C. elegans Lin-7
Mm.2400	glutathione peroxidase 4	10		type 2
Mm.2397	synaptophysin		Mm.203939	expressed sequence Al256814
Mm.23826	phosphotyrosyl phosphatase activator		Mm.203924	expressed sequence AW259572
Mm.2381	amyloid beta (A4) precursor-like protein		Mm.203921	expressed sequence Al850305
	1		Mm.202728	expressed sequence Al447901
Mm.2338	glutamine synthetase	15	Mm.202696	expressed sequence AA409221
Mm.2338	glutamine synthetase		Mm.201729	expressed sequence Al426007
Mm.2326	macrophage migration inhibitory factor		Mm.2011	glutathione S-transferase, mu 1
Mm.2319	Scgn10 like-protein		Mm.200858	RIKEN cDNA 2410129E14 gene
Mm.23023	RIKEN cDNA 1500009C09 gene		Mm.200843	synuclein, beta
Mm.23002	RIKEN cDNA 5330410G16 gene	20	Mm.200817	expressed sequence AW124717
Mm.22699	selenoprotein P, plasma, 1		Mm.200817	expressed sequence AW124717
Mm.22637	RIKEN cDNA 0910001L24 gene		Mm.200806	(Manual) no clear assignment, probably
Mm.22597	RIKEN cDNA 2310042E05 gene			non-coding (but spliced) RNA gene
Mm.22473	Rab acceptor 1 (prenylated)		Mm.200511	expressed sequence Al115024
Mm.22149	succinate dehydrogenase complex,	25	Mm.199903	expressed sequence Al850290
	subunit A, flavoprotein (Fp)		Mm.199652	expressed sequence Al838505
Mm.2214	septin 4		Mm.198588	expressed sequence Al851970
Mm.220966	reticulon 4		Mm.19834	RIKEN cDNA 0610033L03 gene
Mm.220898	calmodulin 3		Mm.197523	brain acyl-CoA hydrolase
Mm.220885	neurochondrin	30	Mm.196614	eukaryotic translation elongation factor
Mm.2206	NADH dehydrogenase (ubiquinone) fla-			1 alpha 1
	voprotein 2		Mm.196611	synapsin I
Mm.219776	RIKEN cDNA 1110001E17 gene		Mm.196607	eukaryotic translation initiation factor 5A
Mm.218848	RIKEN cDNA 3010002G01 gene		Mm.196605	hexokinase 1
Mm.218764	guanine nucleotide binding protein 13,	35	Mm.196578	mitochondrial carrier homolog 1
M 04 004 4	gamma		Mm.196344	lusterin
Mm.218611	receptor (calcitonin) activity modifying		Mm.196239	
M 01740	protein 2			RIKEN cDNA 4922501H04 gene
Mm.21743			Mm.195869	ATPase, H+ transporting, lysosomal
Mm.216438	malate dehydrogenase, mitochondrial	40		ATPase, H+ transporting, lysosomal 31kDa, V1 subunit E
	Mus musculus, clone IMAGE:5068657,	40	Mm.1956	ATPase, H+ transporting, lysosomal 31kDa, V1 subunit E neurofilament, light polypeptide
Mm 216240	Mus musculus, clone IMAGE:5068657, mRNA, partial cds	40		ATPase, H+ transporting, lysosomal 31kDa, V1 subunit E neurofilament, light polypeptide ATP synthase, H+ transporting, mi-
Mm.216240	Mus musculus, clone IMAGE:5068657, mRNA, partial cds Mus musculus, clone IMAGE:3594799,	40	Mm.1956 Mm.19370	ATPase, H+ transporting, lysosomal 31kDa, V1 subunit E neurofilament, light polypeptide ATP synthase, H+ transporting, mitochondrial F1F0 complex, subunit e
	Mus musculus, clone IMAGE:5068657, mRNA, partial cds Mus musculus, clone IMAGE:3594799, mRNA	40	Mm.1956 Mm.19370 Mm.193539	ATPase, H+ transporting, lysosomal 31kDa, V1 subunit E neurofilament, light polypeptide ATP synthase, H+ transporting, mitochondrial F1F0 complex, subunit e H1 histone family, member 2
Mm.21485	Mus musculus, clone IMAGE:5068657, mRNA, partial cds Mus musculus, clone IMAGE:3594799, mRNA RIKEN cDNA 2610102M01 gene		Mm.1956 Mm.19370	ATPase, H+ transporting, lysosomal 31kDa, V1 subunit E neurofilament, light polypeptide ATP synthase, H+ transporting, mitochondrial F1F0 complex, subunit e H1 histone family, member 2 Mus musculus, Similar to metallot-
	Mus musculus, clone IMAGE:5068657, mRNA, partial cds Mus musculus, clone IMAGE:3594799, mRNA RIKEN cDNA 2610102M01 gene Mus musculus, Similar to vesicle-asso-	40 45	Mm.1956 Mm.19370 Mm.193539	ATPase, H+ transporting, lysosomal 31kDa, V1 subunit E neurofilament, light polypeptide ATP synthase, H+ transporting, mitochondrial F1F0 complex, subunit e H1 histone family, member 2 Mus musculus, Similar to metallothionein 1, clone MGC:27821 IMAGE:
Mm.21485	Mus musculus, clone IMAGE:5068657, mRNA, partial cds Mus musculus, clone IMAGE:3594799, mRNA RIKEN cDNA 2610102M01 gene Mus musculus, Similar to vesicle-asso- ciated calmodulin-binding protein, clone		Mm.1956 Mm.19370 Mm.193539 Mm.192991	ATPase, H+ transporting, lysosomal 31kDa, V1 subunit E neurofilament, light polypeptide ATP synthase, H+ transporting, mitochondrial F1F0 complex, subunit e H1 histone family, member 2 Mus musculus, Similar to metallothionein 1, clone MGC:27821 IMAGE: 3483861, mRNA, complete cds
Mm.21485	Mus musculus, clone IMAGE:5068657, mRNA, partial cds Mus musculus, clone IMAGE:3594799, mRNA RIKEN cDNA 2610102M01 gene Mus musculus, Similar to vesicle-associated calmodulin-binding protein, clone MGC:28873 IMAGE:4527857, mRNA,		Mm.1956 Mm.19370 Mm.193539	ATPase, H+ transporting, lysosomal 31kDa, V1 subunit E neurofilament, light polypeptide ATP synthase, H+ transporting, mitochondrial F1F0 complex, subunit e H1 histone family, member 2 Mus musculus, Similar to metallothionein 1, clone MGC:27821 IMAGE: 3483861, mRNA, complete cds amyloid beta (A4) precursor-like protein
Mm.21485 Mm.214549	Mus musculus, clone IMAGE:5068657, mRNA, partial cds Mus musculus, clone IMAGE:3594799, mRNA RIKEN cDNA 2610102M01 gene Mus musculus, Similar to vesicle-associated calmodulin-binding protein, clone MGC:28873 IMAGE:4527857, mRNA, complete cds		Mm.1956 Mm.19370 Mm.193539 Mm.192991 Mm.19133	ATPase, H+ transporting, lysosomal 31kDa, V1 subunit E neurofilament, light polypeptide ATP synthase, H+ transporting, mitochondrial F1F0 complex, subunit e H1 histone family, member 2 Mus musculus, Similar to metallothionein 1, clone MGC:27821 IMAGE: 3483861, mRNA, complete cds amyloid beta (A4) precursor-like protein 2
Mm.21485 Mm.214549 Mm.2133	Mus musculus, clone IMAGE:5068657, mRNA, partial cds Mus musculus, clone IMAGE:3594799, mRNA RIKEN cDNA 2610102M01 gene Mus musculus, Similar to vesicle-associated calmodulin-binding protein, clone MGC:28873 IMAGE:4527857, mRNA, complete cds centaurin, gamma 3	45	Mm.1956 Mm.19370 Mm.193539 Mm.192991 Mm.19133 Mm.19047	ATPase, H+ transporting, lysosomal 31kDa, V1 subunit E neurofilament, light polypeptide ATP synthase, H+ transporting, mitochondrial F1F0 complex, subunit e H1 histone family, member 2 Mus musculus, Similar to metallothionein 1, clone MGC:27821 IMAGE: 3483861, mRNA, complete cds amyloid beta (A4) precursor-like protein 2 expressed sequence Al425998
Mm.21485 Mm.214549 Mm.2133 Mm.212672	Mus musculus, clone IMAGE:5068657, mRNA, partial cds Mus musculus, clone IMAGE:3594799, mRNA RIKEN cDNA 2610102M01 gene Mus musculus, Similar to vesicle-associated calmodulin-binding protein, clone MGC:28873 IMAGE:4527857, mRNA, complete cds centaurin, gamma 3 S100 protein, beta polypeptide, neural		Mm.1956 Mm.19370 Mm.193539 Mm.192991 Mm.19133	ATPase, H+ transporting, lysosomal 31kDa, V1 subunit E neurofilament, light polypeptide ATP synthase, H+ transporting, mitochondrial F1F0 complex, subunit e H1 histone family, member 2 Mus musculus, Similar to metallothionein 1, clone MGC:27821 IMAGE: 3483861, mRNA, complete cds amyloid beta (A4) precursor-like protein 2 expressed sequence Al425998 growth hormone inducible transmem-
Mm.21485 Mm.214549 Mm.2133 Mm.212672 Mm.212516	Mus musculus, clone IMAGE:5068657, mRNA, partial cds Mus musculus, clone IMAGE:3594799, mRNA RIKEN cDNA 2610102M01 gene Mus musculus, Similar to vesicle-associated calmodulin-binding protein, clone MGC:28873 IMAGE:4527857, mRNA, complete cds centaurin, gamma 3 S100 protein, beta polypeptide, neural RIKEN cDNA 2900002L20 gene	45	Mm.1956 Mm.19370 Mm.193539 Mm.192991 Mm.19133 Mm.19047 Mm.182912	ATPase, H+ transporting, lysosomal 31kDa, V1 subunit E neurofilament, light polypeptide ATP synthase, H+ transporting, mitochondrial F1F0 complex, subunit e H1 histone family, member 2 Mus musculus, Similar to metallothionein 1, clone MGC:27821 IMAGE: 3483861, mRNA, complete cds amyloid beta (A4) precursor-like protein 2 expressed sequence Al425998 growth hormone inducible transmembrane protein
Mm.21485 Mm.214549 Mm.2133 Mm.212672	Mus musculus, clone IMAGE:5068657, mRNA, partial cds Mus musculus, clone IMAGE:3594799, mRNA RIKEN cDNA 2610102M01 gene Mus musculus, Similar to vesicle-associated calmodulin-binding protein, clone MGC:28873 IMAGE:4527857, mRNA, complete cds centaurin, gamma 3 S100 protein, beta polypeptide, neural RIKEN cDNA 2900002L20 gene deleted in polyposis 1	45	Mm.1956 Mm.19370 Mm.193539 Mm.192991 Mm.19133 Mm.19047	ATPase, H+ transporting, lysosomal 31kDa, V1 subunit E neurofilament, light polypeptide ATP synthase, H+ transporting, mitochondrial F1F0 complex, subunit e H1 histone family, member 2 Mus musculus, Similar to metallothionein 1, clone MGC:27821 IMAGE: 3483861, mRNA, complete cds amyloid beta (A4) precursor-like protein 2 expressed sequence Al425998 growth hormone inducible transmembrane protein ganglioside-induced differentiation-as-
Mm.21485 Mm.214549 Mm.2133 Mm.212672 Mm.212516 Mm.21251	Mus musculus, clone IMAGE:5068657, mRNA, partial cds Mus musculus, clone IMAGE:3594799, mRNA RIKEN cDNA 2610102M01 gene Mus musculus, Similar to vesicle-associated calmodulin-binding protein, clone MGC:28873 IMAGE:4527857, mRNA, complete cds centaurin, gamma 3 S100 protein, beta polypeptide, neural RIKEN cDNA 2900002L20 gene	45	Mm.1956 Mm.19370 Mm.193539 Mm.192991 Mm.19133 Mm.19047 Mm.182912 Mm.18218	ATPase, H+ transporting, lysosomal 31kDa, V1 subunit E neurofilament, light polypeptide ATP synthase, H+ transporting, mitochondrial F1F0 complex, subunit e H1 histone family, member 2 Mus musculus, Similar to metallothionein 1, clone MGC:27821 IMAGE: 3483861, mRNA, complete cds amyloid beta (A4) precursor-like protein 2 expressed sequence Al425998 growth hormone inducible transmembrane protein ganglioside-induced differentiation-associated-protein 1
Mm.21485 Mm.214549 Mm.2133 Mm.212672 Mm.212516 Mm.21251	Mus musculus, clone IMAGE:5068657, mRNA, partial cds Mus musculus, clone IMAGE:3594799, mRNA RIKEN cDNA 2610102M01 gene Mus musculus, Similar to vesicle-associated calmodulin-binding protein, clone MGC:28873 IMAGE:4527857, mRNA, complete cds centaurin, gamma 3 S100 protein, beta polypeptide, neural RIKEN cDNA 2900002L20 gene deleted in polyposis 1 genes associated with retinoid-IFN-in-	45	Mm.1956 Mm.19370 Mm.193539 Mm.192991 Mm.19133 Mm.19047 Mm.182912 Mm.18218 Mm.181894	ATPase, H+ transporting, lysosomal 31kDa, V1 subunit E neurofilament, light polypeptide ATP synthase, H+ transporting, mitochondrial F1F0 complex, subunit e H1 histone family, member 2 Mus musculus, Similar to metallothionein 1, clone MGC:27821 IMAGE: 3483861, mRNA, complete cds amyloid beta (A4) precursor-like protein 2 expressed sequence Al425998 growth hormone inducible transmembrane protein ganglioside-induced differentiation-associated-protein 1 RIKEN cDNA 2900092E17 gene
Mm.21485 Mm.214549 Mm.2133 Mm.212672 Mm.212516 Mm.21251 Mm.21162	Mus musculus, clone IMAGE:5068657, mRNA, partial cds Mus musculus, clone IMAGE:3594799, mRNA RIKEN cDNA 2610102M01 gene Mus musculus, Similar to vesicle-associated calmodulin-binding protein, clone MGC:28873 IMAGE:4527857, mRNA, complete cds centaurin, gamma 3 S100 protein, beta polypeptide, neural RIKEN cDNA 2900002L20 gene deleted in polyposis 1 genes associated with retinoid-IFN-induced mortality 19	45	Mm.1956 Mm.19370 Mm.193539 Mm.192991 Mm.19133 Mm.19047 Mm.182912 Mm.18218	ATPase, H+ transporting, lysosomal 31kDa, V1 subunit E neurofilament, light polypeptide ATP synthase, H+ transporting, mitochondrial F1F0 complex, subunit e H1 histone family, member 2 Mus musculus, Similar to metallothionein 1, clone MGC:27821 IMAGE: 3483861, mRNA, complete cds amyloid beta (A4) precursor-like protein 2 expressed sequence Al425998 growth hormone inducible transmembrane protein ganglioside-induced differentiation-associated-protein 1 RIKEN cDNA 2900092E17 gene RIKEN cDNA 2610041P16 gene
Mm.21485 Mm.214549 Mm.2133 Mm.212672 Mm.212516 Mm.21251 Mm.21162 Mm.2108	Mus musculus, clone IMAGE:5068657, mRNA, partial cds Mus musculus, clone IMAGE:3594799, mRNA RIKEN cDNA 2610102M01 gene Mus musculus, Similar to vesicle-associated calmodulin-binding protein, clone MGC:28873 IMAGE:4527857, mRNA, complete cds centaurin, gamma 3 S100 protein, beta polypeptide, neural RIKEN cDNA 2900002L20 gene deleted in polyposis 1 genes associated with retinoid-IFN-induced mortality 19 transthyretin	45	Mm.1956 Mm.19370 Mm.193539 Mm.192991 Mm.19133 Mm.19047 Mm.182912 Mm.18218	ATPase, H+ transporting, lysosomal 31kDa, V1 subunit E neurofilament, light polypeptide ATP synthase, H+ transporting, mitochondrial F1F0 complex, subunit e H1 histone family, member 2 Mus musculus, Similar to metallothionein 1, clone MGC:27821 IMAGE: 3483861, mRNA, complete cds amyloid beta (A4) precursor-like protein 2 expressed sequence Al425998 growth hormone inducible transmembrane protein ganglioside-induced differentiation-associated-protein 1 RIKEN cDNA 2900092E17 gene RIKEN cDNA 2610041P16 gene cytochrome c oxidase, subunit Vb
Mm.21485 Mm.214549 Mm.2133 Mm.212672 Mm.212516 Mm.21251 Mm.21162 Mm.2108 Mm.21071	Mus musculus, clone IMAGE:5068657, mRNA, partial cds Mus musculus, clone IMAGE:3594799, mRNA RIKEN cDNA 2610102M01 gene Mus musculus, Similar to vesicle-associated calmodulin-binding protein, clone MGC:28873 IMAGE:4527857, mRNA, complete cds centaurin, gamma 3 S100 protein, beta polypeptide, neural RIKEN cDNA 2900002L20 gene deleted in polyposis 1 genes associated with retinoid-IFN-induced mortality 19 transthyretin ADP-ribosylation-like 2	45	Mm.1956 Mm.19370 Mm.193539 Mm.192991 Mm.19133 Mm.19047 Mm.182912 Mm.18218 Mm.181894 Mm.181721 Mm.180182	ATPase, H+ transporting, lysosomal 31kDa, V1 subunit E neurofilament, light polypeptide ATP synthase, H+ transporting, mitochondrial F1F0 complex, subunit e H1 histone family, member 2 Mus musculus, Similar to metallothionein 1, clone MGC:27821 IMAGE: 3483861, mRNA, complete cds amyloid beta (A4) precursor-like protein 2 expressed sequence Al425998 growth hormone inducible transmembrane protein ganglioside-induced differentiation-associated-protein 1 RIKEN cDNA 2900092E17 gene RIKEN cDNA 2610041P16 gene

17	EP 1 529 838 A1
17	EF 1 323 030 A1

Mm.177117	Mus musculus, clone MGC:31632 IMAGE:4511454, mRNA, complete cds		Mm.115124 Mm.114810	brain protein 14 expressed sequence AW060990
Mm.176927	RIKEN cDNA 2610301115 gene		Mm.1147	Mus musculus calmodulin III (Calm3)
Mm.17484 Mm.16831	synuclein, alpha	-	M 40707	mRNA, 3' untranslated region
Mm.16769	creatine kinase, brain RIKEN cDNA 5031406P05 gene	5	Mm.10727	ATPase, H+ transporting, lysosomal
Mm.16767	heterogeneous nuclear ribonucleopro-		Mm 102700	56/58kD, V1 subunit B, isoform 2
141111.10707	tein A2/B1		Mm.103709	potassium inwardly-rectifying channel,
Mm.16763	aldolase 1, A isoform		Mm.103605	subfamily J, member 10 DnaJ (Hsp40) homolog, subfamily B,
Mm.16228	solute carrier family 25 (mitochondrial	10	Wil 71. 1 0 3 0 0 3	member 10
	carrier; adenine nucleotide transloca-		Mm.102278	secretory carrier membrane protein 5
	tor), member 4		Mm.102276	expressed sequence R74975
Mm.16080	dynamin		Mm.101476	(Manual assignment) BNPI, VGLUT-1,
Mm.158871	RIKEN cDNA 2410003L22 gene			mouse homolog of putative vesicular
Mm.157929	ESTs, Weakly similar to PBAS MOUSE	15		glutamate transporter, Na+/Phosphate
	PROBASIN PRECURSOR			cotransporter
Mm.157859	ESTs		Mm.100980	calneuron 1
Mm.157648	RIKEN cDNA 5730403B10 gene		Mm.1008	prostaglandin D2 synthase (21 kDa,
Mm.15711	cyclic nucleotide phosphodiesterase 1			brain)
Mm.156959	beta-spectrin 4	20	Mm.1008	(Manual) Prostaglandin H2 D-Isomera-
Mm.15571	amyloid beta (A4) precursor protein			se (PGD2 SYNTHASE) (PGDS2)
Mm.15512	potassium voltage-gated channel, sha-			(PGDS) member of lipocalin family
14	ker-related subfamily, beta member 2			
Mm.154651	purine rich element binding protein B		Liste D: Posit	ivmarker neurale Stammzellen (1.);
Mm.153758	RIKEN cDNA 0610040H15 gene	25		
Mm.15125 Mm.14798	stromal cell derived factor receptor 1		ES-Zeilen +; I	PSA-NCAM - ; Adult brain -
Mm.142511	ribosomal protein S13		[0040]	
Mm.142317	expressed sequence Al173355 RIKEN cDNA 2610009E16 gene		[0042]	
Mm.142140	neurofilament, medium polypeptide	30	Mm.9703	(Manual) conner transport pretain/ebs
Mm.140761	DnaJ (Hsp40) homolog, subfamily C,	00	WIII.9703	(Manual) copper transport protein/cha- perone ATOX1
	member 5		Mm.930	cathepsin L
Mm.139797	expressed sequence Al848587		Mm.90787	nerve growth factor receptor
Mm.139239	RIKEN cDNA 2900016C05 gene			(TNFRSF16) associated protein 1
Mm.139239	RIKEN cDNA 2900016C05 gene	35	Mm.90587	enolase 1, alpha non-neuron
Mm.139239	RIKEN cDNA 2900016C05 gene		Mm.90115	lysophospholipase 1
Mm.138866	apolipoprotein E		Mm.90003	gap junction membrane channel prote-
Mm.13859	ribosomal protein L41			in beta 3
Mm.1383	Rho GDP dissociation inhibitor (GDI)		Mm.88302	EST, Weakly similar to S14234 hypo-
	gamma	40		thetical protein - mouse
Mm.135621	expressed sequence Al848120		Mm.88212	tubulin, alpha 6
Mm.13445	3-oxoacid CoA transferase		Mm.87581	(Manual) fibronectin 1, internal tag (ma-
Mm.1339 Mm.131127	chromogranin B		11 277000	jor tag probably AAAAAAAAAA)
Mm.12958	RIKEN cDNA 6230410L23 gene	45	Mm.87293	WD repeat domain 12
Mm.12860	kinesin light chain 2 G protein-coupled receptor 37-like 1	43	Mm.87216	Rab geranylgeranyl transferase, a sub-
Mm.1268	proteolipid protein (myelin)		Mm.8155	unit TG interacting factor
Mm.1268	(Manual assignment) PLP Myelin Pro-		Mm.78861	nucleolar and coiled-body phosphopro-
	teolipid Protein, uh05d10.r1 Soares			tein 1
	mouse hypothalamus NMHy Mus mus-	50	Mm.76780	ESTs
	culus cDNA clone 1617043 5' similar to		Mm.7417	cyclin D3
	gb:M54927 MYELIN PROTEOLIPID		Mm.7387	RNA polymerase 1-4 (194 kDa subunit)
	PROTEIN		Mm.7381	hypoxia induced gene 1
Mm.12468	glioblastoma amplified sequence		Mm.725	ribosomal protein L7a
Mm.124592	expressed sequence AW214631	55	Mm.71046	ESTs
Mm.1239	glial fibrillary acidic protein		Mm.70127	ribosomal protein L12
Mm.1222	brain abundant, membrane attached si-		Mm.69647	pancreas specific transcription factor,
	gnal protein 2			1a

EΡ	1	529	838	A1	

Mm.69049	cDNA sequence AF155546		M 0045	IMAGE:3992883, mRNA, complete cds
Mm.6700	eukaryotic translation initiation factor 4E binding protein 1		Mm.3845	Mus musculus, eukaryotic translation termination factor 1, clone MGC:18745
Mm.66	ribosomal protein S4, X-linked			IMAGE:3992883, mRNA, complete cds
Mm.6579	centromere autoantigen A	5	Mm.38151	adenylosuccinate lyase
Mm.6534 Mm.6343	calpain, small subunit 1		Mm.38057	ESTs
Mm.584	nucleophosmin 1		Mm.3776	Mus musculus, clone MGC:37810
Mm.57223	annexin A2		M- 0750	IMAGE:5098241, mRNA, complete cds
Mm.57153	helicase, lymphoid specific sterol O-acyltransferase 2	10	Mm.3752 Mm.36241	RAN binding protein 1
Mm.5624	DEAD/H (Asp-Glu-Ala-Asp/His) box	10	Mm.360	B-cell receptor-associated protein 37
141111.5024	polypeptide 16		Mm.3572	cytochrome c oxidase, subunit Va RIKEN cDNA 1110033J19 gene
Mm.548	cytochrome c oxidase, subunit VIc		Mm.35621	ESTs
Mm.5305	(Manual) GNB2L1, RACK1, Receptor		Mm.35605	cadherin 1
	of activated C kinase, WD40-repeat	15	Mm.3487	ribosomal protein L30
	protein		Mm.3486	ribosomal protein L3
Mm.5290	(Manual) 60S ribosomal protein L17		Mm.34828	heat shock protein, 105 kDa
	(L23) (popey3-annotation wrong)		Mm.34797	cellular retinoic acid binding protein I
Mm.4993	matrix metalloproteinase 3		Mm.34606	RIKEN cDNA 2610511F02 gene
Mm.493	CCCTC-binding factor	20	Mm.34554	Mus musculus, Similar to E2F trans-
Mm.4890	Finkel-Biskis-Reilly murine sarcoma vi-			cription factor 4, p107/p130-binding,
	rus (FBR-MuSV) ubiquitously expres-			clone MGC:37558 IMAGE:4987691,
	sed (fox derived)			mRNA, complete cds
Mm.4770	frizzled homolog 7 (Drosophila)		Mm.3438	lamin A
Mm.4742	proliferation-associated 2G4, 38kD	25	Mm.34351	Mus musculus, Similar to hypothetical
Mm.46461	L-threonine dehydrogenase			protein FLJ13187, clone MGC:28979
Mm.4606	branched chain aminotransferase 1,			IMAGE:4503757, mRNA, complete cds
Mm 4560	cytosolic		Mm.34102	ornithine decarboxylase, structural
Mm.4560	low density lipoprotein receptor-related protein associated protein 1	30	Mm.3379	serine hydroxymethyl transferase 1
Mm.45237	RIKEN cDNA 2610318N02 gene	30	Mm 22040	(soluble)
Mm.45151	RIKEN cDNA 1700043E15 gene		Mm.33240 Mm.33202	epithelial V-like antigen RIKEN cDNA 2410018A17 gene
Mm.4502	mini chromosome maintenance de-		Mm.32879	testis expressed gene 17
	ficient (S. cerevisiae)		Mm.321	secreted phosphoprotein 1
Mm.43831	lectin, galactose binding, soluble 1	35	Mm.318	RIKEN cDNA 2010107E04 gene
Mm.43162	RIKEN cDNA 0710008D09 gene		Mm.31227	expressed sequence AW123847
Mm.42960	RIKEN cDNA 2610301D06 gene		Mm.30929	peroxiredoxin 1
Mm.4280	RIKEN cDNA 2010203J19 gene		Mm.3049	CDC28 protein kinase 1
Mm.42790	ribosomal protein S18		Mm.30242	peptidylprolyl isomerase D (cyclophilin
Mm.42767	ribosomal protein S17	40		D)
Mm.42197	proteasome (prosome, macropain)		Mm.30184	RIKEN cDNA 2700086123 gene
	subunit, beta type 1		Mm.30114	amyotrophic lateral sclerosis 2 (juveni-
Mm.42196	nuclear protein 95			le) homolog (human)
Mm.42195	RuvB-like protein 1		Mm.30060	RIKEN cDNA 2310008N12 gene
Mm.41467	Mus musculus, clone MGC:28892	45	Mm.30049	complement component 1, q subcom-
Mm 41151	IMAGE:4912251, mRNA, complete cds		14 00004	ponent binding protein
Mm.41151 Mm.41061	ESTS		Mm.30034	translocase of inner mitochondrial
Mm.41	RIKEN cDNA 1810055P05 gene (Manual) Mitochondrial ATP synthase		Mm 20004	membrane 8 homolog a (yeast)
WIIII4 1	oligomycin sensitivity conferral protein	50	Mm.29904 Mm.29902	mitochondrial ribosomal protein L15 Mus musculus, Similar to phosphoseri-
	(OSCP) (ATP5O)		14/111.233UZ	ne aminotransferase, clone MGC:6462
Mm.4095	inactive X specific transcripts			IMAGE:2616298, mRNA, complete cds
Mm.4024	cofilin 1, non-muscle		Mm.29859	eukaryotic translation initiation factor 2,
Mm.3925	S100 calcium binding protein A4			subunit 2 (beta, 38kDa)
Mm.38718	ESTs, Moderately similar to S12207 hy-	55	Mm.29856	RIKEN cDNA 9130022B02 gene
	pothetical protein		Mm.29717	3-monooxgenase/tryptophan 5-mo-
Mm.3845	Mus musculus, eukaryotic translation			nooxgenase activation protein, gamma
	termination factor 1, clone MGC: 18745			polypeptide

EP 1 529 838 A1

Mm.29714	(Manual) mouse version of muscle-			tor), member 13
	specific protein M9		Mm.24506	Mus musculus, clone IMAGE:3591061,
Mm.29675	thioredoxin-like 2			mRNA, partial cds
Mm.29619	RIKEN cDNA 1200007E24 gene		Mm.2437	BING4 protein
Mm.29513	NADH dehydrogenase (ubiquinone) 1	5	Mm.2424	ribosomal protein L10A
	alpha subcomplex, 7 (14.5kD, B14.5a)		Mm.24220	RIKEN cDNA 2310003F16 gene
Mm.29504	sperm specific antigen 1		Mm.24219	RIKEN cDNA 1810037117 gene
Mm.2942	asparagine synthetase		Mm.24174	Mus musculus, similar to alanyl-tRNA
Mm.29405	ring-box 1			synthetase (H. sapiens), clone MGC:
Mm.29363	RIKEN cDNA 2310044F10 gene	10		37368 IMAGE:4976684, mRNA, com-
Mm.2930	Mus musculus, Similar to peter pan			plete cds
	(Drosophila) homolog, clone MGC:		Mm.2395	male enhanced antigen 1
	25669 IMAGE:4489442, mRNA, com-		Mm.2355	prohibitin
	plete cds		Mm.235	ubiquitin B
Mm.29192	asparaginyl-tRNA synthetase	15	Mm.22731	integrin beta 4 binding protein
Mm.29148	RIKEN cDNA 2400008B06 gene		Mm.22626	Mus musculus, Similar to chromosome
Mm.29122	RIKEN cDNA 0610012D09 gene			14 open reading frame 3, clone MGC:
Mm.29076	RIKEN cDNA 2510010F10 gene			36589 IMAGE:5320590, mRNA, com-
Mm.28919	destrin			plete cds
Mm.28892	expressed sequence AA959742	20	Mm.2246	proteasome (prosome, macropain)
Mm.28805	SET translocation			subunit, beta type 7
Mm.2849	heat shock protein, 74 kDa, A		Mm.22421	telomerase binding protein, p23
Mm.28483	Mus musculus, Similar to hypothetical		Mm.22421	telomerase binding protein, p23
	protein FLJ22479, clone IMAGE:		Mm.22317	RIKEN cDNA 8430410A17 gene
	4487274, mRNA, partial cds	25	Mm.22288	cyclin D1
Mm.28405	serum/glucocorticoid regulated kinase		Mm.22271	smt3-specific isopeptidase 1
Mm.28173	ESTs, Moderately similar to JC5224		Mm.220992	Mus musculus, clone IMAGE:3492506,
	methioninetRNA ligase			mRNA, partial cds
Mm.28053	RIKEN cDNA 1110017C15 gene		Mm.219671	Mus musculus, clone MGC:36369
Mm.28035	ESTs, Weakly similar to	30		IMAGE:4982239, mRNA, complete cds
M 07004	TRHY_HUMAN TRICHOHYALI		Mm.219458	RNA binding protein gene with multiple
Mm.27901	RIKEN cDNA 1110020J08 gene			splicing
Mm.27858	RIKEN cDNA 1110036B12 gene		Mm.218533	RIKEN cDNA 1500016H10 gene
Mm.27855	replication factor C (activator 1) 2	-	Mm.2180	heat shock protein, 84 kDa 1
Mm 0750	(40kD)	<i>3</i> 5	Mm.21758	cytochrome P450, 2e1, ethanol induci-
Mm.2758	makorin, ring finger protein, 3		14 . 04000	ble
Mm.27536	ESTs, Highly similar to hypothetical		Mm.21630	expressed sequence AU022237
Mm 27526	protein FLJ14075		Mm.21569	RIKEN cDNA 2700069E09 gene
Mm.27526	(Manual) Arginyl tRNA synthetase (RI-	40	Mm.213020	(Manual) 60S ribosomal protein L32
Mm.27186	KEN cDNA 2610011N19)	40	M 040000	(RPL32)
WIIII.27 100	Mus musculus, Similar to CG7083 ge-		Mm.212899	Mus musculus, Similar to RIKEN cDNA
	ne product, clone MGC:6480 IMAGE: 2646515, mRNA, complete cds			1200009K13 gene, clone MGC: 18794
Mm.2718	eukaryotic translation elongation factor		Mm.21289	IMAGE:4193513, mRNA, complete cds
141111.27 10	1 beta 2	45		ribosomal protein S12
Mm.2718	eukaryotic translation elongation factor	43	Mm.21086	eukaryotic translation elongation factor
141111.27 10	1 beta 2			1 delta (guanine nucleotide exchange
Mm.27134	RIKEN cDNA 2610033C09 gene		Mm.210638	protein) EST
Mm.265	ribosomal protein S25		Mm.210638	
Mm.2647	profilin 1	50	Mm.21054	expressed sequence C87860 nuclease sensitive element binding
Mm.2623	serine (or cysteine) proteinase inhibitor,		WIII1.21054	protein 1
	clade B (ovalbumin), member 6		Mm.20943	FK506 binding protein 9
Mm.25642	RIKEN cDNA 2310034K10 gene		Mm.20925	G1 to phase transition 1
Mm.254	tumor protein, translationally-controlled		Mm.20923	nuclear localization signal protein ab-
	1	55		sent in velo-cardio-facial patients
Mm.25328	ESTs		Mm.20848	regulatory factor X-associated ankyrin-
Mm.24513	solute carrier family 25 (mitochondrial			containing protein
	carrier; adenine nucleotide transloca-		Mm.20847	sorting nexin 5
				-

_	^
,	

Mm.20294	selenophosphate synthetase 2		Mm 157770	DIVEN aDNA 0010004E10 ann
Mm.20290	expressed sequence AW536573		Mm.157778	RIKEN cDNA 2610034E13 gene
Mm.20288	glutathione reductase 1		Mm.154915	ribosomal protein S29
Mm.200920	ribosomal protein S28		Mm.154387	transketolase
Mm.197601	•	-	Mm.153963	CD8 antigen, beta chain
WIIII. 197001	heat shock 10 kDa protein 1 (chapero- nin 10)	5	Mm.153159 Mm.152291	chaperonin subunit 6a (zeta) EST
Mm.197555	hypothetical protein MGC6664			
Mm. 197551	heat shock 70kD protein 8		Mm.151329	karyopherin (importin) beta 3
Mm.196604	angio-associated migratory protein, re-		Mm.148973	RIKEN cDNA 3010025E17 gene
WIIII. 130004		40	Mm.147946	MYB binding protein (P160) 1a
Mm.196586	lated sequence cullin 2	10	Mm.147693	ribosomal protein S3
Mm. 196581	· · · · · -		Mm.14768	reduced expression 3
Mm.196526	mitogen activated protein kinase 1		Mm.14663	ATP synthase, H+ transporting, mi-
	ADP-ribosylation factor 6			tochondrial F0 complex, subunit g
Mm.196396	tubulin, alpha 1		Mm.143141	eukaryotic translation initiation factor
Mm.196081	peptidylprolyl isomerase (cyclophilin)-	15		1A
14 400	like 1		Mm.142740	metallothionein 2
Mm.196	neural precursor cell expressed, de-		Mm.14245	ribosomal protein, large P2
	velopmentally down-regulated gene 8		Mm.14244	ribosomal protein L9
Mm.195894	Mus musculus, clone MGC:11792		Mm.141443	lactate dehydrogenase 1, A chain
	IMAGE:3595167, mRNA, complete cds	20	Mm.141187	trans-golgi network protein 2
Mm.19169	thioredoxin-like (32kD)		Mm.140380	ribosomal protein L23
Mm.188	(Manual) X-linked phosphoglycerate ki-		Mm.139825	Mus musculus, Similar to xylosylprotein
	nase (PGK1)			betal,4-galactosyltransferase, poly-
Mm.18637	teratocarcinoma expressed, serine rich			peptide 7 (galactosyltransferase I), clo-
Mm. 18459	fibroblast growth factor inducible 14	25		ne MGC: 28643 IMAGE:4224150, mR-
Mm.183022	DNA segment, Chr 8, Brigham & Wo-			NA, complete cds
	men's Genetics 1112 expressed		Mm.13705	(Manual) mouse version of p180 ribo-
Mm.182951	proteasome (prosome, macropain) subunit, alpha type 2			some receptor/ribosome binding prote- in 1 RRBP1
Mm.182931	phosphoribosylaminoimidazole car-	30	Mm.13020	ribosomal protein L13a
	boxylase, phosphoribosylaminoribosy-		Mm.12909	amyloid beta (A4) precursor protein-
	laminoimidazole, succinocarboxamide			binding, family A, member 3
	synthetase		Mm.1275	thioredoxin 1
Mm.182471	RIKEN cDNA 2610524G07 gene		Mm.12508	karyopherin (importin) alpha 2
Mm.181765	Mus musculus 8 days embryo whole	35	Mm.1164	SEC61, gamma subunit (S. cerevisiae)
	body cDNA, RIKEN full-length enriched	-	Mm.11376	ribosomal protein L36
	library, clone:5730409M10:CCAAT/en-		Mm.1125	expressed in non-metastatic cells 2,
	hancer binding protein alpha (C/EBP),		WIIII. 1 123	protein (NM23B) (nucleoside diphos-
	related sequence 1, full insert se-			phate kinase)
	quence	40	Mm.1120	
Mm.181740	interferon-related developmental regu-	40	Mm.108076	endometrial bleeding associated factor
	lator 2		Mm.10706	phosphofructokinase, platelet RIKEN cDNA 2010004J23 gene
Mm.180299	DNA segment, Chr 16, Wayne State		Mm.10706	(Manual) mouse version of 60S riboso-
	University 109, expressed			mal protein L4
Mm.17932	purine-nucleoside phosphorylase	45	Mm.10702	calcyclin binding protein
Mm.1777	heat shock protein, 60 kDa		Mm.10665	Mus musculus, clone IMAGE:3498496,
Mm.176845	RIKEN cDNA 1110069M14 gene			mRNA, partial cds
Mm.175848	(Manual) small Ca-binding protein Cal-		Mm.10623	expressed sequence Al480570
	gizzarin (S100A11) (ENDOTHELIAL		Mm.10600	glutamate dehydrogenase
	MONOCYTE-ACTIVATING POLYPEP-	50	Mm.1056	solute carrier family 1, member 7
	TIDE) (EMAP)		Mm.10474	RIKEN cDNA 3110005M08 gene
Mm.175661	RIKEN cDNA 1110036C17 gene		Mm.101619	EST
Mm.1710	hydroxymethylbilane synthase		Mm.10	spermidine synthase
Mm.17031	POU domain, class 5, transcription fac-		Mm.4325	Kruppel-like factor 4 (gut) [Swissprot:
	tor 1	55		splQ60793;splQ9R255;]
Mm.16757	solute carrier family 20, member 1		Mm.12919	insulin-like growth factor 2, binding pro-
Mm.1639	myeloid cell leukemia sequence 1			tein 1 [Swissprot: splO88477;]
Mm.16110	cyclin E1		Mm.20348	nidogen 2 [Swissprot: splO88322;

EP 1	529	838 A	11
		000 /	• •

	splQ8R5G0;splQ9CT94;]		Mm.7793	protein phosphatase 1, catalytic subu-
Mm.34407	MAD homolog 7 (Drosophila) [Swiss-		141111.7733	nit, gamma isoform
	prot: splO35253;splQ9CSC7;]		Mm.7723	poly(A) binding protein, nuclear 1
Mm.4451	hairy and enhancer of split 1, (Droso-		Mm.76278	RIKEN cDNA 2610203K23 gene
	phila) [Swissprot: none]	5	Mm.7516	nuclear autoantigenic sperm protein (hi-
Mm.57195	nodal [Swissprot: spIP43021;]			stone-binding)
Mm.1249	laminin, gamma 1 [Swissprot: spl P02468;]		Mm.7312	DNA segment, Chr 17, human D6S56E
Mm.27706	ash2 (absent, small, or homeotic)-like		Mm.7141	proliferating cell nuclear antigen
	(Drosophila) [Swissprot:	10	Mm.6787	splicing factor, arginine/serine-rich 3
Mm.4603	scavenger receptor class B1 [Swiss-			(SRp20)
	prot: splQ61009;splQ9CWJ7;]		Mm.66	ribosomal protein S4, X-linked
Mm.181562	adhesion regulating molecule 1 [Swis-		Mm.6476	RIKEN cDNA 2700084L22 gene
	sprot: splQ8VCl8;splQ922A7;		Mm.64104	RIKEN cDNA 2410016F19 gene
Mm,43444	splQ9JKV1;]	15	Mm.6343	nucleophosmin 1
MIII.40444	MAD2 (mitotic arrest deficient, homolog)-like 1 (yeast) [Swissprot:		Mm.61901	expressed sequence AI429604
Mm.103675	sacsin [Swissprot: none]		Mm.6065 Mm.5624	inosine 5'-phosphate dehydrogenase 2 DEAD/H (Asp-Glu-Ala-Asp/His) box po-
Mm.980	tenascin C [Swissprot: splQ64706;		141111111111111111111111111111111111111	lypeptide 16
	splQ9WUU4;]	20	Mm.548	cytochrome c oxidase, subunit VIc
Mm.5090	cripto, teratocarcinoma-derived growth		Mm.5305	guanine nucleotide binding protein, beta
	factor (Tdgf1)			2, related sequence 1
Mm.30177	D11Ertd603e, DNA segment, Chr 11,		Mm.525	eukaryotic translation initiation factor 4,
	ERATO Doi 603			gamma 2
Mm.233844	C330012H03Rik, RIKEN cDNA	25	Mm.5114	dishevelled 2, dsh homolog (Drosophi-
	C330012H03		M 4000	la)
Liste F: Posit	tivmarker neurale Stammzellen (2.);		Mm.4933	mini chromosome maintenance de- ficient 6 (S. cerevisiae)
2.010 2. 1 00.	avindraci iledizie otdimizenen (2.),		Mm.4890	Finkel-Biskis-Reilly murine sarcoma vi-
			10000	Time Dens Helly Hallie Salcolla VI-
ES-Zellen +;	PSA-NCAM -/+; Adult brain -	30		rus (FBR-MuSV) ubiquitously expres-
ES-Zellen +;	PSA-NCAM -/+; Adult brain -	30		rus (FBR-MuSV) ubiquitously expressed (fox derived)
ES-Zellen +;	PSA-NCAM -/+; Adult brain -	30	Mm.4846	
[0043]		30	Mm.4846 Mm.4756	sed (fox derived)
[0043] Mm.99776	cytosolic aminopeptidase P		Mm.4756 Mm.46754	sed (fox derived) lamin B1 leptin receptor expressed sequence Al316867
[0043] Mm.99776 Mm.9916	cytosolic aminopeptidase P RIKEN cDNA 1200008012 gene	35	Mm.4756 Mm.46754 Mm.46533	sed (fox derived) lamin B1 leptin receptor expressed sequence Al316867 RIKEN cDNA 1110007L15 gene
[0043] Mm.99776 Mm.9916 Mm.99	cytosolic aminopeptidase P RIKEN cDNA 1200008012 gene ribonucleotide reductase M2		Mm.4756 Mm.46754 Mm.46533 Mm.4551	sed (fox derived) lamin B1 leptin receptor expressed sequence Al316867 RIKEN cDNA 1110007L15 gene villin 2
[0043] Mm.99776 Mm.9916 Mm.99 Mm.9811	cytosolic aminopeptidase P RIKEN cDNA 1200008012 gene ribonucleotide reductase M2 RIKEN cDNA 2310008M10 gene		Mm.4756 Mm.46754 Mm.46533	sed (fox derived) lamin B1 leptin receptor expressed sequence Al316867 RIKEN cDNA 1110007L15 gene villin 2 ATPase, Na+/K+ transporting, beta 1
[0043] Mm.99776 Mm.9916 Mm.99	cytosolic aminopeptidase P RIKEN cDNA 1200008012 gene ribonucleotide reductase M2 RIKEN cDNA 2310008M10 gene (Manual) uncharacterized protein corre-		Mm.4756 Mm.46754 Mm.46533 Mm.4551 Mm.4550	sed (fox derived) lamin B1 leptin receptor expressed sequence Al316867 RIKEN cDNA 1110007L15 gene villin 2 ATPase, Na+/K+ transporting, beta 1 polypeptide
[0043] Mm.99776 Mm.9916 Mm.99 Mm.9811	cytosolic aminopeptidase P RIKEN cDNA 1200008012 gene ribonucleotide reductase M2 RIKEN cDNA 2310008M10 gene		Mm.4756 Mm.46754 Mm.46533 Mm.4551	sed (fox derived) lamin B1 leptin receptor expressed sequence Al316867 RIKEN cDNA 1110007L15 gene villin 2 ATPase, Na+/K+ transporting, beta 1 polypeptide SRY-box containing gene 2
[0043] Mm.99776 Mm.9916 Mm.99 Mm.9811	cytosolic aminopeptidase P RIKEN cDNA 1200008012 gene ribonucleotide reductase M2 RIKEN cDNA 2310008M10 gene (Manual) uncharacterized protein corre- sponding to human splQ9Y3l0, similar	3 5	Mm.4756 Mm.46754 Mm.46533 Mm.4551 Mm.4550	sed (fox derived) lamin B1 leptin receptor expressed sequence Al316867 RIKEN cDNA 1110007L15 gene villin 2 ATPase, Na+/K+ transporting, beta 1 polypeptide
[0043] Mm.99776 Mm.9916 Mm.99 Mm.9811 Mm.9257	cytosolic aminopeptidase P RIKEN cDNA 1200008012 gene ribonucleotide reductase M2 RIKEN cDNA 2310008M10 gene (Manual) uncharacterized protein corre- sponding to human splQ9Y3I0, similar to E.coli rtcB, UPF0027-family transcription factor Dp 1 heat shock 70kD protein 5 (glucose-re-	3 5	Mm.4756 Mm.46754 Mm.46533 Mm.4551 Mm.4550 Mm.4541 Mm.45312	sed (fox derived) lamin B1 leptin receptor expressed sequence Al316867 RIKEN cDNA 1110007L15 gene villin 2 ATPase, Na+/K+ transporting, beta 1 polypeptide SRY-box containing gene 2 anaphase-promoting complex subunit 5
[0043] Mm.99776 Mm.9916 Mm.99 Mm.9811 Mm.9257 Mm.925 Mm.918	cytosolic aminopeptidase P RIKEN cDNA 1200008012 gene ribonucleotide reductase M2 RIKEN cDNA 2310008M10 gene (Manual) uncharacterized protein corre- sponding to human splQ9Y3I0, similar to E.coli rtcB, UPF0027-family transcription factor Dp 1 heat shock 70kD protein 5 (glucose-re- gulated protein, 78kD)	3 5	Mm.4756 Mm.46754 Mm.46533 Mm.4551 Mm.4550 Mm.4541 Mm.45312 Mm.45149 Mm.45132 Mm.4426	sed (fox derived) lamin B1 leptin receptor expressed sequence Al316867 RIKEN cDNA 1110007L15 gene villin 2 ATPase, Na+/K+ transporting, beta 1 polypeptide SRY-box containing gene 2 anaphase-promoting complex subunit 5 ESTs expressed sequence AW121759 Cd63 antigen
[0043] Mm.99776 Mm.9916 Mm.99 Mm.9811 Mm.9257	cytosolic aminopeptidase P RIKEN cDNA 1200008012 gene ribonucleotide reductase M2 RIKEN cDNA 2310008M10 gene (Manual) uncharacterized protein corre- sponding to human splQ9Y3I0, similar to E.coli rtcB, UPF0027-family transcription factor Dp 1 heat shock 70kD protein 5 (glucose-re- gulated protein, 78kD) high mobility group nucleosomal bin-	<i>35</i>	Mm.4756 Mm.46754 Mm.46533 Mm.4551 Mm.4550 Mm.4541 Mm.45312 Mm.45149 Mm.45132	sed (fox derived) lamin B1 leptin receptor expressed sequence Al316867 RIKEN cDNA 1110007L15 gene villin 2 ATPase, Na+/K+ transporting, beta 1 polypeptide SRY-box containing gene 2 anaphase-promoting complex subunit 5 ESTs expressed sequence AW121759 Cd63 antigen MAD2 (mitotic arrest deficient, homo-
[0043] Mm.99776 Mm.9916 Mm.99 Mm.9811 Mm.9257 Mm.925 Mm.918 Mm.911	cytosolic aminopeptidase P RIKEN cDNA 1200008012 gene ribonucleotide reductase M2 RIKEN cDNA 2310008M10 gene (Manual) uncharacterized protein corre- sponding to human splQ9Y3I0, similar to E.coli rtcB, UPF0027-family transcription factor Dp 1 heat shock 70kD protein 5 (glucose-re- gulated protein, 78kD) high mobility group nucleosomal bin- ding domain 2	3 5	Mm.4756 Mm.46754 Mm.46533 Mm.4551 Mm.4550 Mm.4541 Mm.45312 Mm.45149 Mm.45132 Mm.4426 Mm.43444	sed (fox derived) lamin B1 leptin receptor expressed sequence Al316867 RIKEN cDNA 1110007L15 gene villin 2 ATPase, Na+/K+ transporting, beta 1 polypeptide SRY-box containing gene 2 anaphase-promoting complex subunit 5 ESTs expressed sequence AW121759 Cd63 antigen MAD2 (mitotic arrest deficient, homolog)-like 1 (yeast)
[0043] Mm.99776 Mm.9916 Mm.99 Mm.9811 Mm.9257 Mm.925 Mm.918	cytosolic aminopeptidase P RIKEN cDNA 1200008012 gene ribonucleotide reductase M2 RIKEN cDNA 2310008M10 gene (Manual) uncharacterized protein corre- sponding to human splQ9Y3I0, similar to E.coli rtcB, UPF0027-family transcription factor Dp 1 heat shock 70kD protein 5 (glucose-re- gulated protein, 78kD) high mobility group nucleosomal bin- ding domain 2 heterogeneous nuclear ribonucleopro-	<i>35</i>	Mm.4756 Mm.46754 Mm.46533 Mm.4551 Mm.4550 Mm.4541 Mm.45312 Mm.45149 Mm.45132 Mm.4426 Mm.43444 Mm.4280	sed (fox derived) lamin B1 leptin receptor expressed sequence Al316867 RIKEN cDNA 1110007L15 gene villin 2 ATPase, Na+/K+ transporting, beta 1 polypeptide SRY-box containing gene 2 anaphase-promoting complex subunit 5 ESTs expressed sequence AW121759 Cd63 antigen MAD2 (mitotic arrest deficient, homolog)-like 1 (yeast) RIKEN cDNA 2010203J19 gene
[0043] Mm.99776 Mm.9916 Mm.99 Mm.9811 Mm.9257 Mm.925 Mm.918 Mm.911	cytosolic aminopeptidase P RIKEN cDNA 1200008012 gene ribonucleotide reductase M2 RIKEN cDNA 2310008M10 gene (Manual) uncharacterized protein corre- sponding to human splQ9Y3I0, similar to E.coli rtcB, UPF0027-family transcription factor Dp 1 heat shock 70kD protein 5 (glucose-re- gulated protein, 78kD) high mobility group nucleosomal bin- ding domain 2 heterogeneous nuclear ribonucleopro- tein L	<i>35</i>	Mm.4756 Mm.46754 Mm.46533 Mm.4551 Mm.4550 Mm.4541 Mm.45312 Mm.45132 Mm.45132 Mm.4426 Mm.43444 Mm.4280 Mm.42767	sed (fox derived) lamin B1 leptin receptor expressed sequence Al316867 RIKEN cDNA 1110007L15 gene villin 2 ATPase, Na+/K+ transporting, beta 1 polypeptide SRY-box containing gene 2 anaphase-promoting complex subunit 5 ESTs expressed sequence AW121759 Cd63 antigen MAD2 (mitotic arrest deficient, homolog)-like 1 (yeast) RIKEN cDNA 2010203J19 gene ribosomal protein S17
[0043] Mm.99776 Mm.9916 Mm.99 Mm.9811 Mm.9257 Mm.925 Mm.918 Mm.911 Mm.9043	cytosolic aminopeptidase P RIKEN cDNA 1200008012 gene ribonucleotide reductase M2 RIKEN cDNA 2310008M10 gene (Manual) uncharacterized protein corre- sponding to human splQ9Y3I0, similar to E.coli rtcB, UPF0027-family transcription factor Dp 1 heat shock 70kD protein 5 (glucose-re- gulated protein, 78kD) high mobility group nucleosomal bin- ding domain 2 heterogeneous nuclear ribonucleopro-	<i>35</i>	Mm.4756 Mm.46754 Mm.46533 Mm.4551 Mm.4550 Mm.4541 Mm.45312 Mm.45132 Mm.45132 Mm.4426 Mm.43444 Mm.4280 Mm.42767 Mm.4237	sed (fox derived) lamin B1 leptin receptor expressed sequence Al316867 RIKEN cDNA 1110007L15 gene villin 2 ATPase, Na+/K+ transporting, beta 1 polypeptide SRY-box containing gene 2 anaphase-promoting complex subunit 5 ESTs expressed sequence AW121759 Cd63 antigen MAD2 (mitotic arrest deficient, homolog)-like 1 (yeast) RIKEN cDNA 2010203J19 gene ribosomal protein S17 topoisomerase (DNA) II alpha
[0043] Mm.99776 Mm.9916 Mm.99 Mm.9811 Mm.9257 Mm.925 Mm.918 Mm.911 Mm.9043 Mm.89927	cytosolic aminopeptidase P RIKEN cDNA 1200008012 gene ribonucleotide reductase M2 RIKEN cDNA 2310008M10 gene (Manual) uncharacterized protein corre- sponding to human splQ9Y3I0, similar to E.coli rtcB, UPF0027-family transcription factor Dp 1 heat shock 70kD protein 5 (glucose-re- gulated protein, 78kD) high mobility group nucleosomal bin- ding domain 2 heterogeneous nuclear ribonucleopro- tein L signal recognition particle 9 kDa	<i>35</i>	Mm.4756 Mm.46754 Mm.46533 Mm.4551 Mm.4550 Mm.4541 Mm.45312 Mm.45132 Mm.45132 Mm.4426 Mm.43444 Mm.4280 Mm.42767	sed (fox derived) lamin B1 leptin receptor expressed sequence Al316867 RIKEN cDNA 1110007L15 gene villin 2 ATPase, Na+/K+ transporting, beta 1 polypeptide SRY-box containing gene 2 anaphase-promoting complex subunit 5 ESTs expressed sequence AW121759 Cd63 antigen MAD2 (mitotic arrest deficient, homolog)-like 1 (yeast) RIKEN cDNA 2010203J19 gene ribosomal protein S17
[0043] Mm.99776 Mm.9916 Mm.99 Mm.9811 Mm.9257 Mm.925 Mm.918 Mm.911 Mm.9043 Mm.89927 Mm.89579	cytosolic aminopeptidase P RIKEN cDNA 1200008012 gene ribonucleotide reductase M2 RIKEN cDNA 2310008M10 gene (Manual) uncharacterized protein corre- sponding to human splQ9Y3I0, similar to E.coli rtcB, UPF0027-family transcription factor Dp 1 heat shock 70kD protein 5 (glucose-re- gulated protein, 78kD) high mobility group nucleosomal bin- ding domain 2 heterogeneous nuclear ribonucleopro- tein L signal recognition particle 9 kDa mannose-P-dolichol utilization defect 1 H3 histone, family 3A tubulin, alpha 6	35 40 45	Mm.4756 Mm.46754 Mm.46533 Mm.4551 Mm.4550 Mm.4541 Mm.45312 Mm.45132 Mm.45132 Mm.4426 Mm.43444 Mm.4280 Mm.42767 Mm.4237	sed (fox derived) lamin B1 leptin receptor expressed sequence Al316867 RIKEN cDNA 1110007L15 gene villin 2 ATPase, Na+/K+ transporting, beta 1 polypeptide SRY-box containing gene 2 anaphase-promoting complex subunit 5 ESTs expressed sequence AW121759 Cd63 antigen MAD2 (mitotic arrest deficient, homolog)-like 1 (yeast) RIKEN cDNA 2010203J19 gene ribosomal protein S17 topoisomerase (DNA) II alpha proteasome (prosome, macropain) sub-
[0043] Mm.99776 Mm.9916 Mm.99 Mm.9811 Mm.9257 Mm.925 Mm.918 Mm.911 Mm.9043 Mm.89927 Mm.89579 Mm.89136 Mm.88212 Mm.880	cytosolic aminopeptidase P RIKEN cDNA 1200008012 gene ribonucleotide reductase M2 RIKEN cDNA 2310008M10 gene (Manual) uncharacterized protein corre- sponding to human splQ9Y3I0, similar to E.coli rtcB, UPF0027-family transcription factor Dp 1 heat shock 70kD protein 5 (glucose-re- gulated protein, 78kD) high mobility group nucleosomal bin- ding domain 2 heterogeneous nuclear ribonucleopro- tein L signal recognition particle 9 kDa mannose-P-dolichol utilization defect 1 H3 histone, family 3A tubulin, alpha 6 mammary tumor integration site 6	35 40 45	Mm.4756 Mm.46754 Mm.46754 Mm.46533 Mm.4551 Mm.4550 Mm.4541 Mm.45312 Mm.45149 Mm.45132 Mm.4426 Mm.4280 Mm.42767 Mm.42767 Mm.42197 Mm.4215 Mm.4215 Mm.41940	sed (fox derived) lamin B1 leptin receptor expressed sequence Al316867 RIKEN cDNA 1110007L15 gene villin 2 ATPase, Na+/K+ transporting, beta 1 polypeptide SRY-box containing gene 2 anaphase-promoting complex subunit 5 ESTs expressed sequence AW121759 Cd63 antigen MAD2 (mitotic arrest deficient, homolog)-like 1 (yeast) RIKEN cDNA 2010203J19 gene ribosomal protein S17 topoisomerase (DNA) II alpha proteasome (prosome, macropain) subunit, beta type 1
[0043] Mm.99776 Mm.9916 Mm.99 Mm.9811 Mm.9257 Mm.925 Mm.918 Mm.911 Mm.9043 Mm.89927 Mm.89579 Mm.89579 Mm.88552	cytosolic aminopeptidase P RIKEN cDNA 1200008012 gene ribonucleotide reductase M2 RIKEN cDNA 2310008M10 gene (Manual) uncharacterized protein corre- sponding to human splQ9Y3I0, similar to E.coli rtcB, UPF0027-family transcription factor Dp 1 heat shock 70kD protein 5 (glucose-re- gulated protein, 78kD) high mobility group nucleosomal bin- ding domain 2 heterogeneous nuclear ribonucleopro- tein L signal recognition particle 9 kDa mannose-P-dolichol utilization defect 1 H3 histone, family 3A tubulin, alpha 6 mammary tumor integration site 6 baculoviral IAP repeat-containing 5	35 40 45	Mm.4756 Mm.46754 Mm.46754 Mm.46533 Mm.4551 Mm.4550 Mm.4541 Mm.45312 Mm.45149 Mm.45132 Mm.4426 Mm.4280 Mm.42767 Mm.42767 Mm.42197 Mm.42197	sed (fox derived) lamin B1 leptin receptor expressed sequence Al316867 RIKEN cDNA 1110007L15 gene villin 2 ATPase, Na+/K+ transporting, beta 1 polypeptide SRY-box containing gene 2 anaphase-promoting complex subunit 5 ESTs expressed sequence AW121759 Cd63 antigen MAD2 (mitotic arrest deficient, homolog)-like 1 (yeast) RIKEN cDNA 2010203J19 gene ribosomal protein S17 topoisomerase (DNA) II alpha proteasome (prosome, macropain) subunit, beta type 1 catalase 1 RIKEN cDNA 6530409L22 gene cyclin A2
[0043] Mm.99776 Mm.9916 Mm.99 Mm.9811 Mm.9257 Mm.925 Mm.918 Mm.911 Mm.9043 Mm.89927 Mm.89579 Mm.89136 Mm.88212 Mm.880	cytosolic aminopeptidase P RIKEN cDNA 1200008012 gene ribonucleotide reductase M2 RIKEN cDNA 2310008M10 gene (Manual) uncharacterized protein corre- sponding to human splQ9Y3I0, similar to E.coli rtcB, UPF0027-family transcription factor Dp 1 heat shock 70kD protein 5 (glucose-re- gulated protein, 78kD) high mobility group nucleosomal bin- ding domain 2 heterogeneous nuclear ribonucleopro- tein L signal recognition particle 9 kDa mannose-P-dolichol utilization defect 1 H3 histone, family 3A tubulin, alpha 6 mammary tumor integration site 6 baculoviral IAP repeat-containing 5 KH domain containing, RNA binding, si-	35 40 45	Mm.4756 Mm.46754 Mm.46754 Mm.46533 Mm.4551 Mm.4550 Mm.4541 Mm.45312 Mm.45149 Mm.45132 Mm.4426 Mm.4280 Mm.42767 Mm.42767 Mm.42197 Mm.4215 Mm.41940 Mm.4189 Mm.41023	sed (fox derived) lamin B1 leptin receptor expressed sequence Al316867 RIKEN cDNA 1110007L15 gene villin 2 ATPase, Na+/K+ transporting, beta 1 polypeptide SRY-box containing gene 2 anaphase-promoting complex subunit 5 ESTs expressed sequence AW121759 Cd63 antigen MAD2 (mitotic arrest deficient, homolog)-like 1 (yeast) RIKEN cDNA 2010203J19 gene ribosomal protein S17 topoisomerase (DNA) II alpha proteasome (prosome, macropain) subunit, beta type 1 catalase 1 RIKEN cDNA 6530409L22 gene cyclin A2 RIKEN cDNA 1110021E09 gene
[0043] Mm.99776 Mm.9916 Mm.99 Mm.9811 Mm.9257 Mm.925 Mm.918 Mm.911 Mm.9043 Mm.89927 Mm.89579 Mm.89579 Mm.885212 Mm.880 Mm.8552 Mm.8256	cytosolic aminopeptidase P RIKEN cDNA 1200008012 gene ribonucleotide reductase M2 RIKEN cDNA 2310008M10 gene (Manual) uncharacterized protein corre- sponding to human splQ9Y3I0, similar to E.coli rtcB, UPF0027-family transcription factor Dp 1 heat shock 70kD protein 5 (glucose-re- gulated protein, 78kD) high mobility group nucleosomal bin- ding domain 2 heterogeneous nuclear ribonucleopro- tein L signal recognition particle 9 kDa mannose-P-dolichol utilization defect 1 H3 histone, family 3A tubulin, alpha 6 mammary tumor integration site 6 baculoviral IAP repeat-containing 5 KH domain containing, RNA binding, si- gnal transduction associated 1	35 40 45	Mm.4756 Mm.46754 Mm.46754 Mm.46533 Mm.4551 Mm.4550 Mm.4541 Mm.45312 Mm.45149 Mm.45132 Mm.4426 Mm.4280 Mm.42767 Mm.42767 Mm.42197 Mm.42197	sed (fox derived) lamin B1 leptin receptor expressed sequence Al316867 RIKEN cDNA 1110007L15 gene villin 2 ATPase, Na+/K+ transporting, beta 1 polypeptide SRY-box containing gene 2 anaphase-promoting complex subunit 5 ESTs expressed sequence AW121759 Cd63 antigen MAD2 (mitotic arrest deficient, homolog)-like 1 (yeast) RIKEN cDNA 2010203J19 gene ribosomal protein S17 topoisomerase (DNA) II alpha proteasome (prosome, macropain) subunit, beta type 1 catalase 1 RIKEN cDNA 6530409L22 gene cyclin A2 RIKEN cDNA 1110021E09 gene antigen identified by monoclonal antibo-
[0043] Mm.99776 Mm.9916 Mm.99 Mm.9811 Mm.9257 Mm.925 Mm.918 Mm.911 Mm.9043 Mm.89927 Mm.89579 Mm.89579 Mm.88552	cytosolic aminopeptidase P RIKEN cDNA 1200008012 gene ribonucleotide reductase M2 RIKEN cDNA 2310008M10 gene (Manual) uncharacterized protein corre- sponding to human splQ9Y3I0, similar to E.coli rtcB, UPF0027-family transcription factor Dp 1 heat shock 70kD protein 5 (glucose-re- gulated protein, 78kD) high mobility group nucleosomal bin- ding domain 2 heterogeneous nuclear ribonucleopro- tein L signal recognition particle 9 kDa mannose-P-dolichol utilization defect 1 H3 histone, family 3A tubulin, alpha 6 mammary tumor integration site 6 baculoviral IAP repeat-containing 5 KH domain containing, RNA binding, si- gnal transduction associated 1 TG interacting factor	35 40 45	Mm.4756 Mm.46754 Mm.46754 Mm.46533 Mm.4551 Mm.4550 Mm.4541 Mm.45312 Mm.45132 Mm.45132 Mm.4426 Mm.43444 Mm.4280 Mm.42767 Mm.42767 Mm.42197 Mm.4215 Mm.41940 Mm.4189 Mm.41023 Mm.44078	sed (fox derived) lamin B1 leptin receptor expressed sequence Al316867 RIKEN cDNA 1110007L15 gene villin 2 ATPase, Na+/K+ transporting, beta 1 polypeptide SRY-box containing gene 2 anaphase-promoting complex subunit 5 ESTs expressed sequence AW121759 Cd63 antigen MAD2 (mitotic arrest deficient, homolog)-like 1 (yeast) RIKEN cDNA 2010203J19 gene ribosomal protein S17 topoisomerase (DNA) II alpha proteasome (prosome, macropain) subunit, beta type 1 catalase 1 RIKEN cDNA 6530409L22 gene cyclin A2 RIKEN cDNA 1110021E09 gene antigen identified by monoclonal antibody Ki 67
[0043] Mm.99776 Mm.9916 Mm.99 Mm.9811 Mm.9257 Mm.925 Mm.918 Mm.911 Mm.9043 Mm.89927 Mm.89579 Mm.89579 Mm.885212 Mm.880 Mm.8552 Mm.8256 Mm.8155	cytosolic aminopeptidase P RIKEN cDNA 1200008012 gene ribonucleotide reductase M2 RIKEN cDNA 2310008M10 gene (Manual) uncharacterized protein corre- sponding to human splQ9Y3I0, similar to E.coli rtcB, UPF0027-family transcription factor Dp 1 heat shock 70kD protein 5 (glucose-re- gulated protein, 78kD) high mobility group nucleosomal bin- ding domain 2 heterogeneous nuclear ribonucleopro- tein L signal recognition particle 9 kDa mannose-P-dolichol utilization defect 1 H3 histone, family 3A tubulin, alpha 6 mammary tumor integration site 6 baculoviral IAP repeat-containing 5 KH domain containing, RNA binding, si- gnal transduction associated 1	35 40 45	Mm.4756 Mm.46754 Mm.46754 Mm.46533 Mm.4551 Mm.4550 Mm.4541 Mm.45312 Mm.45149 Mm.45132 Mm.4426 Mm.4280 Mm.42767 Mm.42767 Mm.42197 Mm.4215 Mm.41940 Mm.4189 Mm.41023	sed (fox derived) lamin B1 leptin receptor expressed sequence Al316867 RIKEN cDNA 1110007L15 gene villin 2 ATPase, Na+/K+ transporting, beta 1 polypeptide SRY-box containing gene 2 anaphase-promoting complex subunit 5 ESTs expressed sequence AW121759 Cd63 antigen MAD2 (mitotic arrest deficient, homolog)-like 1 (yeast) RIKEN cDNA 2010203J19 gene ribosomal protein S17 topoisomerase (DNA) II alpha proteasome (prosome, macropain) subunit, beta type 1 catalase 1 RIKEN cDNA 6530409L22 gene cyclin A2 RIKEN cDNA 1110021E09 gene antigen identified by monoclonal antibo-

EP 1	529	838	A1	

Mm 4004	nafilia 4 nan massata			
Mm.4024	cofilin 1, non-muscle		Mm.29122	RIKEN cDNA 0610012D09 gene
Mm.3931	Max protein		Mm.29055	chromobox homolog 1 (Drosophila HP1
Mm.38930	expressed sequence AA407558			beta)
Mm.38912	RIKEN cDNA 2410129H14 gene		Mm.29054	RIKEN cDNA 2610529I12 gene
Mm.38611	RIKEN cDNA 2210021A15 gene	5	Mm.29005	expressed sequence AU021749
Mm.38528	RIKEN cDNA 2810430M08 gene		Mm.28995	RIKEN cDNA 2010009J12 gene
Mm.38306	macrophage erythroblast attacher		Mm.28985	ribosomal protein L27
Mm.3797	nucleosome assembly protein 1-like 1		Mm.28965	RIKEN cDNA 0710007A14 gene
Mm.37835	ribosomal protein L7		Mm.28964	Mus musculus, clone IMAGE:4949762,
Mm.372	ribosomal protein S26	10		mRNA, partial cds
Mm.36511	mitochondrial ribosomal protein L32		Mm.28961	cleavage and polyadenylation specific
Mm.35844	growth arrest specific 5			factor 5, 25 kD subunit
Mm.35829	erythroid differentiation regulator		Mm.28909	protein tyrosine phosphatase 4a1
Mm.35661	Mus musculus, Similar to hypothetical		Mm.28899	RIKEN cDNA 1110059P08 gene
	protein, clone MGC:29235 IMAGE:	15	Mm.28805	SET translocation
	5043282, mRNA, complete cds		Mm.28805	SET translocation
Mm.35087	expressed sequence AA673488		Mm.28805	SET translocation
Mm.3501	kinesin family member C5A		Mm.28726	EST C77032
Mm.34914	ESTs		Mm.28694	RIKEN cDNA 2410088K19 gene
Mm.3487	ribosomal protein L30	20	Mm.28560	Ly1 antibody reactive clone
Mm.3444	bromodomain-containing 2		Mm.28499	Mus musculus, similar to CG15881 ge-
Mm.34385	expressed sequence Al450344			ne product (H. sapiens), clone MGC:
Mm.34261	expressed sequence AW557761			36308 IMAGE:5040108, mRNA, com-
Mm.3381	ribosomal protein S8			plete cds
Mm.3380	kinesin family member 5B	25	Mm.28299	ESTs, Highly similar to GUAA_HUMAN
Mm.3360	tyrosine 3-monooxygenase/tryptophan			GMP SYNTHASE
	5-monooxygenase activation protein,		Mm.28222	RIKEN cDNA 2610307C23 gene
	zeta polypeptide		Mm.28121	RIKEN cDNA 1110061A19 gene
Mm.326	RIKEN cDNA 1110038L14 gene		Mm.28044	filamin-like protein
Mm.320	DNA polymerase alpha 2, 68 kDa	30	Mm.27972	NS1-associated protein 1
Mm.3199	RIKEN cDNA 1500001N04 gene		Mm.27927	heterogeneous nuclear ribonucleopro-
Mm.31512	ring finger protein 2			tein A1
Mm.31228	RIKEN cDNA 1810022K09 gene		Mm.27852	expressed sequence AW555814
Mm.30806	ribosomal protein L19		Mm.27818	eukaryotic translation elongation factor
Mm.3054	alpha-L-iduronidase	35		2
Mm.3035	RIKEN cDNA 3110006P09 gene		Mm.27796	RIKEN cDNA 5730427N09 gene
Mm.30270	proteasome (prosome, macropain) sub-		Mm.27669	small nuclear ribonucleoprotein E
	unit, alpha type 4		Mm.27660	RIKEN cDNA 5730420G12 gene
Mm.30120	ribosomal protein S27-like		Mm.27624	RIKEN cDNA C530002L11 gene
Mm.30069	RIKEN cDNA 1200003J11 gene	40	Mm.27293	RIKEN cDNA 4833420K19 gene
Mm.30011	ribosomal protein S23		Mm.27269	RIKEN cDNA 2310037I24 gene
Mm.29931	cell division cycle 20 homolog (S. cere-		Mm.27141	Rac GTPase-activating protein 1
	visiae)		Mm.27074	RIKEN cDNA 2610019N13 gene
Mm.29923	SMT3 (supressor of mif two, 3) homolog		Mm.265	ribosomal protein S25
	2 (S. cerevisiae)	45	Mm.2591	RNA binding motif protein 3
Mm.29911	RIKEN cDNA 3200001M24 gene		Mm.25558	RIKEN cDNA 2410018J24 gene
Mm.29896	ribosomal protein L21		Mm.25542	(Manual) strange EST contig in intron of
Mm.2986	expressed sequence AW146116			p137 (GPI-anchored transcytosis prote-
Mm.29829	expressed sequence Al326010			in), maybe alternative C-terminus of
Mm.29666	solute carrier family 25 (mitochondrial	50		splQ60865
	carnitine/acylcarnitine translocase),		Mm.254	tumor protein, translationally-controlled
	member 20			1
Mm.2966	isocitrate dehydrogenase 2 (NADP+),		Mm.25299	ESTs, Weakly similar to simple repeat
	mitochondrial			sequence-containing transcript
Mm.29296	RIKEN cDNA 1110003H02 gene	55	Mm.25164	gene trap locus 1-13
Mm.29194	RIKEN cDNA 1700094M07 gene		Mm.25137	RIKEN cDNA 2410004B18 gene
Mm.29133	budding uninhibited by benzimidazoles		Mm.24870	(Manual assignment) UBP7 ubiquitin
	1 homolog, beta (S. cerevisiae)			hydrolase

ΕP	1	529	838	A 1	

Mm.24591 Mm.2424	expressed sequence AW546279 ribosomal protein L10A		Mm.21054	nuclease sensitive element binding protein 1
Mm.24219 Mm.24042	RIKEN cDNA 1810037I17 gene RIKEN cDNA 1210001E11 gene	_	Mm.20927	transforming growth factor beta 1 indu- ced transcript 4
Mm.23943 Mm.23758	vesicle-associated membrane protein, associated protein A (33 kDa) RIKEN cDNA 1110008P04 gene	5	Mm.206399 Mm.2038	ESTs Ras-GTPase-activating protein SH3- domain binding protein
Mm.23695	dihydrofolate reductase		Mm.2025	survival motor neuron
Mm.23692	casein kinase II, alpha 1 related sequence 4	10	Mm.200837	Mus musculus, clone IMAGE:5355658, mRNA
Mm.23096	protein phosphatase 2 (formerly 2A), regulatory subunit B", alpha		Mm.196614	eukaryotic translation elongation factor 1 alpha 1
Mm.2287	proteasome (prosome, macropain) sub- unit, alpha type 5		Mm.196608 Mm.196526	expressed sequence AA407306 ADP-ribosylation factor 6
Mm.22731	integrin beta 4 binding protein	15	Mm.196515	DNA segment, Chr 1, ERATO Doi 692,
Mm.2265	U1 small nuclear ribonucleoprotein 1C			expressed
Mm.22387	expressed sequence Al314668		Mm.196396	tubulin, alpha 1
Mm.22269	exportin 1, CRM1 homolog (yeast)		Mm.196365	RIKEN cDNA 4833416109 gene
Mm.22214	RIKEN cDNA 2610008F03 gene		Mm.196328	RIKEN cDNA 5830466J11 gene
Mm.220918	heterogeneous nuclear ribonucleoprotein D-like	20	Mm.195898	phosphatidylethanolamine binding protein
Mm.220342	Mus musculus, clone IMAGE:3669867,		Mm.1951	ribonucleic acid binding protein S1
	mRNA, partial cds		Mm.1948	t-complex testis expressed 1
Mm.219670	Mus musculus, Similar to eukaryotic		Mm.193688	RIKEN cDNA 2700059D21 gene
	translation initiation factor 4 gamma, 1,	25	Mm.19187	prothymosin alpha
	clone IMAGE:4950789, mRNA, partial cds		Mm.19101	DEAD (aspartate-glutamate-alanine-
Mm.219668	RIKEN cDNA 2610209F03 gene		Mm.19015	aspartate) box polypeptide 5 serine racemase
Mm.219648	Mus musculus, Similar to nuclear matrix		Mm.18923	mini chromosome maintenance de-
	protein p84, clone MGC:28284 IMAGE:	30	WIIII. 10923	ficient 7 (S. cerevisiae)
	4010605, mRNA, complete cds		Mm.18921	valosin containing protein
Mm.21964	Mus musculus, clone IMAGE:3485208,		Mm. 18856	mitogen-activated protein kinase 6
	mRNA, partial cds		Mm.18705	vacuolar protein sorting 4b (yeast)
Mm.21873	retroviral integration site 1		Mm.18700	RIKEN cDNA 1200009K13 gene
Mm.218657	cerebellar ataxia 3	35	Mm.18637	teratocarcinoma expressed, serine rich
Mm.21841	splicing factor, arginine/serine-rich 2		Mm.18516	H3 histone, family 3B
Mm.218240	(SC-35) Mus musculus, clone IMAGE:5342828,		Mm.1843 Mm.183102	heat shock protein, 86 kDa 1
141111.210240	mRNA, partial cds		Mm.183016	actin-related protein 3 homolog (yeast) thymine DNA glycosylase
Mm.2180	heat shock protein, 84 kDa 1	40	Mm.181880	RIKEN cDNA 1110007A14 gene
Mm.21764	small nuclear ribonucleoprotein poly-		Mm.181562	adhesion regulating molecule 1
	peptide G		Mm.1815	cytidine 5'-triphosphate synthase
Mm.21714	RIKEN cDNA 2410003A14 gene		Mm.180873	RIKEN cDNA 2510019J09 gene
Mm.21559	non-POU-domain-containing, octamer		Mm.180873	(Manual) probably reverse tag of 60S ri-
	binding protein	45		bosomal protein L18a
Mm.213452	Mus musculus, clone IMAGE:5320271, mRNA, partial cds		Mm.180409 Mm.180271	ubiquitin-conjugating enzyme E2H RIKEN cDNA 5630400D24 gene
Mm.213020	(Manual) 60S ribosomal protein L32 (RPL32)		Mm.17989 Mm.1777	chaperonin subunit 8 (theta) heat shock protein, 60 kDa
Mm.21295	expressed sequence AW214031	50	Mm.1775	hematological and neurological expres-
Mm.21289	ribosomal protein S12			sed sequence 1
Mm.21281	ring finger protein 4		Mm.177451	RIKEN cDNA 5730544L10 gene
Mm.21185	adaptor-related protein complex AP-3,		Mm.17330	ESTs
Mm.2115	beta 1 subunit heterogeneous nuclear ribonucleopro-	55	Mm.17306	tropomyosin 3, gamma
···III.2113	tein U	"	Mm.1703 Mm.16976	tubulin, beta 5 TAF9 RNA polymerase II, TATA box bin-
Mm.21094	DNA segment, Chr 9, Wayne State Uni-		WIII. 103/0	ding protein (TBP)-associated factor, 32
	versity 138, expressed			kDa
	·			

= 1	1 5	29	838

Mm.16775 ribosomal protein S24 Mm.16767 heterogeneous nuclear ribonucleoprotein A2/B1 Mm.16711 mini chromosome maintenance deficient 2 (S. cerevisiae) 5 Mm.16525 polo-like kinase homolog, (Drosophila) Mm.1639 myeloid cell leukemia sequence 1 Mm.16323 eukaryotic translation initiation factor 4A2 Mm.16323 eukaryotic translation initiation factor 10 4A2 Mm.156892 heterogeneous nuclear ribonucleoprotein D Mm.15571 amyloid beta (A4) precursor protein Mm.154915 ribosomal protein S29 15 Mm.153457 RIKEN cDNA 2810406C15 gene Mm.148973 RIKEN cDNA 3010025E17 gene Mm.142872 heterogeneous nuclear ribonucleoprotein K Mm.14245 ribosomal protein, large P2 20 Mm.14244 ribosomal protein L9 Mm.142363 RIKEN cDNA 2810036L13 gene Mm.140804 Mus musculus, guanine nucleotide binding protein (G protein), gamma 5, clone MGC:8292 IMAGE:3593324, mRNA, complete cds Mm.140380 ribosomal protein L23 Mm.13886 suppressor of initiator codon mutations, related sequence 1 (S. cerevisiae) Mm.133825 RIKEN cDNA 0610010123 gene 30 Mm.13356 RIKEN cDNA 3110079L04 gene Mm.131705 Mus musculus, Similar to single-stranded DNA binding protein, clone MGC: 41439 IMAGE: 1314987, mRNA, complete cds 35 Mm.12858 eukaryotic translation initiation factor Mm.12706 Mus musculus, Similar to CG11246 gene product, clone MGC:8248 IMAGE: 3591968, mRNA, complete cds 40 Mm.12604 sirtuin 1 ((silent mating type information regulation 2, homolog) 1 (S. cerevisiae) Mm.12568 expressed sequence AW541137 Mm.12508 karyopherin (importin) alpha 2 Mm. 12441 expressed sequence AU014645 45 Mm.124 thymopoietin Mm.12236 zinc finger protein 207 Mm.12145 retinoblastoma binding protein 4 Mm.116989 actin-like Mm.111 poly(rC) binding protein 2 50 Mm.10706 RIKEN cDNA 2010004J23 gene Mm.10474 RIKEN cDNA 3110005M08 gene Mm.10409 golgi autoantigen, golgin subfamily a, 4 Mm.103675 sacsin Mm.1013 ligase I, DNA, ATP-dependent 55 Mm. 101274 RIKEN cDNA 2010008E23 gene Mm.10076 mitochondrial ribosomal protein L13 Mm. 16469 Nmycl, neuroblastoma myc-related on-

31

cogene 1

Patentansprüche

A1

 Zellpopulation, dadurch gekennzeichnet, dass mindestens 5% der Zellen neurale Vorläuferzellen sind, die wenigstens einen der in Liste A oder Liste B aufgeführten Marker aufweisen.

32

- Zellpopulation, dadurch gekennzeichnet, dass mindestens 5% der Zellen, neurale Vorläuferzellen sind, die wenigstens zwei, bevorzugt wenigstens 3 der in Liste A oder Liste B aufgeführten Marker aufweisen.
- Zellpopulation, nach mindestens einem der Ansprüche 1 bis 2, dadurch gekennzelchnet, dass die neuralen Vorläuferzellen keinen in Liste C aufgeführten Marker aufweisen.
- Zellpopulation nach mindestens einem der Ansprüche 1 bis 3, dadurch gekennzeichnet, dass mindestens 25 % der Zellen neurale Vorläuferzellen sind.
- Zellpopulation nach mindestes einem der Ansprüche 1 bis 4, dadurch gekennzeichnet, dass es sich um eine murine Zellpopulation handelt und/ oder die neuralen Vorläuferzellen aus Hirngewebe erhältlich ist.
- Verfahren zur Isolierung einer Zellpopulation nach mindestens einem der Ansprüche 1 bis 5 mit folgenden Schritten:
 - a) Entnahme einer Probe aus dem Hirn
 - b) Isolieren der neuralen Vorläuferzellen unter Verwendung der angegebenen Marker

oder

- a) Differenzierung von embryonalen Stammzellen zu neuralen Vorläuferzellen,
- b) Isolieren der neuralen Vorläuferzellen unter Verwendung der angegebenen Marker

oder

- a) Trans-Differenzierung von adulten, nicht neuralen Stammzellen zu neuralen Vorläuferzellen,
- b) Isolieren der neuralen Vorläuferzeilen unter Verwendung der angegebenen Marker

oder

EP 1 529 838 A1

5

10

20

34

- a) Differenzierung von adulten, neuralen Stammzellen zu neuralen Vorläuferzellen.
- b) Isolieren der neuralen Vorläuferzellen unter Verwendung der angegebenen Marker

oder

- a) Differenzierung von immortalisierten Zellen zu neuralen Vorläuferzellen,
- b) Isolieren der neuralen Vorläuferzellen unter Verwendung der angegebenen Marker.
- Verwendung mindestens eines Markers ausgewählt aus der Liste A oder Liste B zu Identifizierung oder Isolierung von neuralen Vorläuferzellen.
- Antikörper gegen einen Marker aus der Liste A, B oder C.
- Diagnostikmittel enthaltend mindestens einen, bevorzugt zwei oder mehr Substanzen zur Erkennung der Marker der Liste A, B oder C.
- Arzneimittel enthaltend die Zellpopulation nach einem der Ansprüche 1 bis 5.
- Zellpopulation, dadurch gekennzeichnet, dass mindestens 5% der Zellen neurale Stammzellen sind, die wenigstens einen der in Liste D oder Liste E aufgeführten Marker aufweisen.
- Zellpopulation, dadurch gekennzeichnet, dass mindestens 5% der Zellen neurale Stammzellen sind, die wenigstens zwei, bevorzugt wenigstens 3 der in Liste D oder Liste E aufgeführten Marker aufweisen.
- 13. Zellpopulation, nach mindestens einem der Ansprüche 11 bis 12, dadurch gekennzeichnet, dass die neuralen Stammzellen keinen in Liste A oder Liste C aufgeführten Marker aufweisen.
- Zellpopulation nach mindestens einem der Ansprüche 11-13, dadurch gekennzeichnet, dass mindestens 25% der Zellen neurale Stammzellen sind.
- 15. Zellpopulation nach mindestes einem der Ansprüche 11 bis 14, dadurch gekennzeichnet, dass es sich um eine murine Zellpopulation handelt und/oder die neuralen Stammzellen aus Hirngewebe erhältlich
- 16. Verfahren zur Isolierung einer Zellpopulation nach mindestens einem der Ansprüche 11 bis 15 mit folgenden Schritten:
 - a) Entnahme einer Probe aus dem Hirn

b) Isolieren der neuralen Stammzellen unter Verwendung der angegebenen Marker

oder

- a) Differenzierung von embryonalen Stammzellen zu neuralen Stammzellen,
- b) Isolieren der neuralen Stammzellen unter Verwendung der angegebenen Marker

oder

- a) Trans-Differenzierung von adulten, nicht neuralen Stammzellen zu neuralen Stammzellen.
- b) Isolieren der neuralen Stammzellen unter Verwendung der angegebenen Marker

oder

- a) De-Differenzierung von adulten, neuralen Vorläuferzellen zu neuralen Stammzellen.
- b) Isolieren der neuralen Stammzellen unter Verwendung der angegebenen Marker

oder

- a) Differenzierung von immortalisierten Zellen zu neuralen Stammzellen,
- b) Isolieren der neuralen Stammzellen unter Verwendung der angegebenen Marker.
- Antikörper gegen einen Marker aus der Liste D, E, A oder C.
- 18. Diagnostikmittel enthaltend mindestens einen, bevorzugt zwei oder mehr Substanzen zur Erkennung der Marker der Liste D, E, A oder C.
- Arzneimittel enthaltend die Zellpopulation nach einem der Ansprüche 11 bis 15.



EP D FORM 1503 00.82 (P04C09)

Europäisches EUROPÄISCHER TEILRECHERCHENBERICHT

Nummer der Anmeldung

der nach Regel 45 des Europäischen Patemübereinkommens für das weitere Verfahren als europäischer Recherchenbericht gilt

EP 03 02 5506

		an optaisoner mooneronenbenent gilt		Ì
	EINSCHLÄGIG	E DOKUMENTE	,	
Kategorie	Kennzeichnung des Doku der maßgebliche	ments mit Angabe, soweit erforderlich en Teile	Betrifft Anspruch	KLASSIFIKATION DER ANMELDUNG (Int.Cl.7)
X	ARSENIJEVIC YVAN E multipotent neural the cortex of the EXPERIMENTAL NEURO Bd. 170, Nr. 1, Ju Seiten 48-62, XP00; ISSN: 0014-4886 * Seite 52, linke rechte Spalte, Abs	T AL: "Isolation of precursors residing in adult human brain" LOGY, li 2001 (2001-07), 2275728 Spalte, letzter Absatz - atz 1 * Spalte, letzter Absatz * Spalte, Absatz 2 -	1-6,10	C12N5/06 G01N33/53 RECHERCHIERTE SACHGEBIETE (Int.Cl.7) C12N G01N
Die Reche in einem s der Techni	LLSTÄNDIGE RECHE rohenabtellung ist der Auffassung, di olohen Umfang nicht entspricht bzw. k für diese Ansprüche nicht, bzw. nus g recherchierte Patentansprüche:	aß ein oder mehrere Ansprüche, den Vorschrifte entsprechen, deß sinnvolle Ermittlungen über d	en des EPÜ len Stand	GOIN
	dig recherchierte Patentansprüche:			
Nicht rech	erchierte Patentansprüche:			
	de Beschränkung der Recherche: e Ergänzungsblatt (:		
	Recherchenort	Abschlußdatum der Recherche	l	Protei
	MÜNCHEN	7. April 2004	Niel	ouhr-Ebel, K
X : von b Y : von b ander A : techn	TEGORIE DER GENANNTEN DOKL sesonderer Bedeutung allein betracht esonderer Bedeutung in Verbindung en Veröffentlichung derselben Kateg ologischer Hintergrund schriffliche Offenbanng	neonen oder Grundsätze h erst ern oder loht worden ist urnent Detreimstimmendes		



EUROPÄISCHER TEILRECHERCHENBERICHT

Nummer der Anmeldung EP 03 02 5506

	EINSCHLÄGIGE DOKUMENTE	KLASSIFIKATION DER ANMELDUNG (Int.CI.7)	
Kategorie	Kennzeichnung des Dokuments mit Angabe, soweit erforderlich der maßgeblichen Teile	Betrifft Anspruch	(industry
	UCHIDA N ET AL: "Direct isolation of human central nervous system stem cells" PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA, NATIONAL ACADEMY OF SCIENCE. WASHINGTON, US, Bd. 97, Nr. 26, 19. Dezember 2000 (2000-12-19), Seiten 14720-14725, XP002223508 ISSN: 0027-8424 * Zusammenfassung * * Seite 14722, rechte Spalte, letzter Absatz - Seite 14724, rechte Spalte, Absatz 1 * * Abbildungen 1,2 *	11-16,19	RECHERCHIERTE
X	KANEKO Y ET AL: "MUSASHI1: AN EVOLUTIONALLY CONSERVED MARKER FOR CNS PROGENITOR CELLS INCLUDING NEURAL STEM CELLS" DEVELOPMENTAL NEUROSCIENCE, S. KARGER, BASEL, CH, Bd. 22, Nr. 1/2, 2000, Seiten 139-153, XP001033925 ISSN: 0378-5866 * Zusammenfassung * * Abbildung 5 *	11-16,19	SACHGEBIETE (Int.cl.)
X	EP 1 354 943 A (NAT INST OF ADVANCED IND SCIEN) 22. Oktober 2003 (2003-10-22) "Monoclonal antibodies, hybridomas, cell isolation method, isolated cells and immunological diagnostic method" * Spalte 2, Zeile 42 - Spalte 3, Zeile 11 * * Spalte 13, Zeile 41 - Spalte 14, Zeile 53 *	1-19	



EUROPÄISCHER TEILRECHERCHENBERICHT

Nummer der Anmeldung EP 03 02 5506

	EINSCHLÄGIGE DOKUMENTE	KLASSIFIKATION DER ANMELDUNG (Int.CL7)	
Kategorie	Kennzeichnung des Dokuments mit Angabe, soweit erforderlich der maßgeblichen Teile	Betrifft Anspruch	
X	GIMONA MARIO ET AL: "Beta-Actin Specific Monoclonal Antibody" CELL MOTILITY AND THE CYTOSKELETON, Bd. 27, Nr. 2, 1994, Seiten 108-116, XP009028901 ISSN: 0886-1544 * das ganze Dokument *	8,9,17, 18	RECHERCHIERTE SACHGEBIETE (Int.Cl.7)
	-		·



UNVOLLSTÄNDIGE RECHERCHE ERGÄNZUNGSBLATT C

Nummer der Anmeldung EP 03 02 5506

Unvollständig recherchierte Ansprüche: 6, 16

Grund für die Beschränkung der Recherche (nicht patentfähige Erfindung(en)):

Artikel 52 (4) EPÜ – Verfahren zur chirurgischen Behandlung des menschlichen oder tierischen Körpers

Weitere Beschränkung der Recherche

Unvollständig recherchierte Ansprüche: 1-5, 7-15, 17-19

Grund für die Beschränkung der Recherche:

In den Listen A-E, auf die sich in den Patentansprüchen bezogen wird, sind insgesamt etwa 1000 putative Positiv- und Negativmarker neuraler Vorläuferzellen und neuraler Stammzellen aufgelistet. Diese putativen Marker sind teilweise bereits bekannte Proteine, wie z.B. beta-Aktin oder Interleukin 1 alpha, teilweise aber auch undefinierte, als "ESTs" benannte sogenannte Marker oder partielle mRNA-Sequenzen. Aufgrund der grossen Anzahl der putativen Marker und deren tw. mangelhaften Identifikation ist es unmöglich, eine vollständige Recherche zu erstellen.

ANHANG ZUM EUROPÄISCHEN RECHERCHENBERICHT ÜBER DIE EUROPÄISCHE PATENTANMELDUNG NR.

EP 03 02 5506

In diesem Anhang sind die Mitglieder der Patentfamilien der im obengenannten europäischen Recherchenbericht angeführten Patentdokumente angegeben.
Die Angaben über die Familienmitglieder entsprechen dem Stand der Datei des Europäischen Patentamts am Diese Angaben dienen nur zur Unterrichtung und erfolgen ohne Gewähr.

07-04-2004

an	lm Recherchenbericht angeführtes Patentdokument		Datum der Veröffentlichung		Mitglied(er) der Patentfamilie	Datum der Veröffentlichung
EP	1354943	Α	22-10-2003	EP JP US	1354943 A2 2004002350 A 2003186335 A1	22-10-2003 08-01-2004 02-10-2003

Für nähere Einzelheiten zu diesem Anhang : siehe Amtsblatt des Europäischen Patentamts, Nr.12/82